

**Time trends in the incidence of work-related ill-health
in the UK, 1996-2014: estimation from THOR
surveillance data**

Report to the UK Health and Safety Executive

**Carder M, McNamee R, Gittins M,
Hussey L, Agius R**

Centre for Occupational and Environmental Health, Centre for Epidemiology, Institute of
Population Health, Faculty of Medical and Human Sciences, the University of Manchester

<http://www.coeh.man.ac.uk/thor>

31st August 2015

KEY MESSAGES

- The incidence of contact dermatitis (CD) as reported by dermatologists to EPIDERM decreased during 1996-2006, after which it remained relatively flat until 2012, and then decreased again between 2012 and 2013. There was little evidence of a further change between 2013 and 2014. The annual average change in CD incidence (1996-2014) was -3.7% (95% CIs: -4.3, -3.2). Analyses of shorter-term trends (2006 to 2014) suggested a similar annual average decrease of -3.9% (95% CIs: -5.5, -2.2) per year.
- Markedly different neoplasia trends were observed depending on whether analyses were based on reports from 'core' or 'sample' dermatologists; the former suggested a decrease in incidence and the latter an increase. Of the two, it is possible 'sample' data are more representative. However, for both groups the confidence intervals on the annual plots were wide and overlapping. It is therefore difficult to draw any firm conclusions about neoplasia trends from these data.
- Overall, the incidence of work-related respiratory disease fell between 1999 and 2007, after which it remained relatively flat. There was slight evidence consistent with a further decrease between 2013 and 2014. The average decrease in asthma incidence (1999-2014) was -7.2% (95% CIs: -8.4, -6.0) per year. Analyses of shorter-term trends (2007 to 2014) showed an average change of -4.4% (95% CIs: -8.1, -0.6) per year.
- Overall, the annual pattern of change for non-malignant pleural disease (NMPD) and mesothelioma incidence was relatively flat. However, for both groups there was evidence of a significant decrease in incidence between 2013 and 2014. For NMPD the average change in incidence was -1.2% (95% CIs: -2.3, -.01) per year whilst for mesothelioma it was -3.0% (95% CIs: -4.3, -1.6). However (especially when considering information from other sources) the results for mesothelioma in particular should be viewed very cautiously as they may reflect a shift in referral patterns rather than a 'true' trend (a proportion of such cases are now referred to oncologists rather than exclusively to chest physicians as used to be the case).
- The data continue to suggest that the incidence of pneumoconiosis has been increasing since (approximately) 2007 with an average increase of +3.2% (95% CIs: +1.1, +5.3) per year (1999-2014). For the period 2007-2014, the equivalent estimate was +10.4% (95% CIs: +4.5, +16.5). There was some suggestion of a further increase between 2013 and 2014 but confidence intervals are wide and overlapping.
- Because of methodological changes, analyses based on cases reported by general practitioners to THOR-GP for the period 2011 onwards (i.e. since the change to 100% 'sample' reporting) are the most informative for current and future trends for this group. These data suggest an average annual decrease in the incidence of total work-related illness of -14.6% (95% CIs: -21.2, -7.5). Equivalent figures for skin were -19.2% (95% CIs: -37.8, +4.9), for musculoskeletal -12.1% (95% CIs: -21.3, -1.7) and for mental ill-health -15% (95% CIs: -25.1, -3.5).
- The estimates provided in this report were not adjusted for the potential impact of reporter 'fatigue'. However, a summary of previous analyses investigating this phenomenon has been provided. Based on these, the likely impact on the average, annual change in incidence for total skin disease (EPIDERM) and total respiratory disease (SWORD) would be to attenuate the estimate from -3.8% to -3.4% and from -3.2% to -2.7%, respectively. There was no evidence of fatigue in THOR-GP sample reporters.

EXECUTIVE SUMMARY

This report describes temporal trends in incidence of work-related illness (WRI) in the UK as reported to the three constituent schemes of THOR which are currently funded by HSE. These are EPIDERM (dermatologists), SWORD (chest physicians) and THOR-GP (GPs). It updates on previously submitted reports by the incorporation of a further year (2014) of data. Data were analysed in a manner (using a 'multi-level' statistical model) in which the number of reported cases over time could be investigated whilst taking into account other factors that might influence the trend, for example, change in the number of physicians reporting or in the number of people employed in the UK. Change in incidence has been presented either as the average, annual percentage change in incidence rate over a defined period or as graphs showing the risk for each year relative to a reference year (2014).

For each scheme, analyses were carried out separately, for the total reported cases and then for each of the conditions of interest (for example, asthma). THOR physicians can participate either on a monthly basis (termed 'core' reporters) or for one randomly allocated month per year (termed 'sample' reporters) and separate analyses were carried out for each of these groups as well as (where appropriate) both types together. Both EPIDERM and SWORD comprise (and have done throughout the study period) a smaller 'core' group (approximately 10% of reporters) and a larger 'sample' group with most physicians remaining as either 'core' or 'sample' throughout their time in the scheme. THOR-GP differs in that all physicians initially participated as 'core' but over time the proportion of 'sample' reporters increased and from 2011 onwards all physicians have been 'sample'. Thus, some physicians changed their reporting frequency from 'core' to 'sample' during their time in the scheme. This is important because evidence (from THOR-GP and the occupational physician reporting scheme, OPRA) has shown that physicians behave differently depending on whether they are participating as 'core' or 'sample' with the former reporting less cases (in any given month) compared to the latter. Because of these extensive changes, GP analyses based on 'sample' reporters only for the period 2011 onwards are probably the most informative of current and future trends for this physician group.

Another important issue to consider is that the longer a physician participates in a voluntary scheme such as THOR they might start to 'fatigue'. This could manifest in a number of ways, some of which might unduly influence the trend. Identifying and adjusting for the potential impact of 'fatigue' has been a difficult methodological

challenge for this project and a summary of previous analyses investigating this phenomenon has been provided. The results of these analyses suggest that, for EPIDERM and SWORD, some of the observed decrease in disease incidence over time is in fact due to reporter 'fatigue' rather than a 'true trend'. Evidence of fatigue was also observed for THOR-GP core reporters but not sample. For EPIDERM, adjusting for 'fatigue' would likely mean (on average) a decrease in (total skin disease) incidence of -3.4% per year (compared to -3.8% per year if not adjusted for). For SWORD (total respiratory disease) the equivalent change would be from -3.2% to -2.7%.

WORK-RELATED SKIN DISEASE: Dermatologists reported a total of 18438 cases of work-related skin disease to EPIDERM (1996-2014), of which 82% were contact dermatitis (CD) and 12% were neoplasia. The annual average decrease in CD incidence (1996-2014) was -3.7% (95% CIs: -4.3, -3.2) and this remained similar when analyses were restricted to 'core' only or 'sample' only cases. The graphs showing relative risk by year (compared to 2014) suggest an initial decrease in incidence in the earlier part of the study period (1996-2006) followed by a relatively flat trend (2006-2012) and a further decrease between 2012 and 2013. There was little suggestion of a further decrease between 2013 and 2014. The average annual, estimated change in CD incidence since 2006 was -3.9% (95% CIs: -5.5, -2.2).

Markedly different neoplasia trends were observed depending on whether analyses were based on case reports from 'core' or 'sample' reporters. The former suggested an annual average decrease in incidence of -4.6% (95% CIs: -6.2, -3.0) whilst the latter suggested an *increase* in incidence of +2.6% (95% CIs: -0.5, +5.7). It is therefore difficult to draw any firm conclusions about skin neoplasia trends from these data. It is possible that 'sample' data are more representative for this disease group ('core' reporters to EPIDERM are a self-selected group of 'keen specialists' whose main area of expertise is likely to be contact dermatitis and therefore other cases, such as neoplasia, may be triaged to other e.g. sample reporters). However, for both groups, the confidence intervals on the annual plots are wide and overlapping. This suggests that dermatologists in general (or those reporting to EPIDERM) are seeing relatively few neoplasia cases and it may be that other physicians, for example oncologists, would be a better source of information about trends in incidence for this disease.

Skin cases are also reported to THOR-GP by general practitioners, comprising 10% of the total cases reported to THOR-GP. The average, annual change in (total skin

disease) incidence based on sample reports only (2011-2014) was -19.2% (95% CIs: -37.8, +4.9).

WORK-RELATED RESPIRATORY DISEASE: Chest physicians reported a total of 12304 cases of work-related respiratory disease to SWORD (1999-2014), of which 19% were asthma, with the remainder being the (primarily) asbestos related diseases; benign pleural plaques (43%), and mesothelioma (19%), as well as pneumoconiosis (9%). Reports from chest physicians suggested that the incidence of asthma decreased during the study period by, on average, -7.2% (95% CIs: -8.4, -6.0) per year (which remained similar when analyses were restricted to 'core' only or 'sample' only cases). The graphs showing relative risk by year (compared to 2014) suggest that asthma incidence initially fell between 1999 and 2007, after which it remained relatively flat with little indication of a significant change between 2013 and 2014. The average annual, estimated change in asthma incidence since 2007 was -4.4% (95% CIs: -8.1, -0.6).

For mesothelioma, the results suggest an average decrease of approximately -3% (95% CIs: -4.3, -1.6) per year. The annual plots show a relatively flat trend since approximately 2010 but there is suggestion of a significant fall in incidence between 2013 and 2014 (which is particularly pronounced in the 'core' reports). For non-malignant pleural disease the average, annual change in incidence was -1.2% (95% CIs: -2.3, -0.1) with some slight variation when analyses were restricted to 'core' data or 'sample' data only. However (especially when considering information from other sources) the results for mesothelioma in particular should be viewed very cautiously as they may reflect a shift in referral patterns rather than a 'true' trend (a proportion of such cases are now referred to oncologists rather than exclusively to chest physicians as used to be the case).

Data from SWORD continue to suggest a possible increase in pneumoconiosis incidence since approximately 2007. The average, annual change (1999-2014) in incidence was +3.2% (95% CIs: +1.1, +5.3) and for 2007-2013 it was +10.4% (95% CIs: +4.5, +16.5).

Analyses based on cases of work-related respiratory disease reported to THOR-GP data were not carried out (only 2% of case reports to THOR-GP are in this category).

WORK-RELATED MUSCULOSKELETAL DISEASE: Musculoskeletal diagnoses reported by GPs to THOR-GP were predominantly (85%) upper limb disorders (hand/wrist/arm/elbow/shoulder) and spine back disorders (neck/thoracic spine/lumbar spine/trunk). Based on cases reported by 'sample' reporters only (2011-2014) an (average) decrease in incidence of total musculoskeletal disorders of -12.1% (95% CIs: -21.3, -1.7) per year was suggested. The annual plots suggest a relatively flat trend between 2011 and 2013 followed by a decrease between 2013 and 2014 (although confidence intervals for the individual year estimates were wide and overlapping). Analyses restricted to ('sample') case reports of spine/back disorders also suggested a decrease in incidence: an average, annual decrease of -17.3% (95% CIs: -30.3, -1.9) with again the largest drop seeming to occur between 2013 and 2014. For the other two diagnostic groups (upper limb and lower limb) the confidence intervals were very wide making it difficult to draw any firm conclusions at this stage.

WORK-RELATED MENTAL ILL-HEALTH: Mental ill-health case reports to THOR-GP were predominantly (96%) other work stress and anxiety and depression. Based on cases reported by 'sample' reporters (2011 to 2014), an (average) annual change in incidence of total mental ill-health of -15% (95% CIs: -25.1, -3.5) was suggested. As observed for musculoskeletal disorders, the largest drop in incidence again appears to have occurred between 2013 and 2014. A decrease in incidence was also suggested for anxiety and depression at -20.6% (95% CIs: -36.6, -0.5) per year and for other work stress at -9.7% (95% CIs: -22.2, +4.7).

EXECUTIVE SUMMARY

CONTENTS

	Page number
1 BACKGROUND	12
2 METHOD	12
2.1 DATA PERIOD	13
2.2 REPORTER GROUPS	13
2.3 CATEGORIES OF ILLNESS	14
2.4 THE MULTI-LEVEL MODEL AND ITS ASSUMPTIONS	15
2.5 SUMMARY OF 'REPORTER FATIGUE' INVESTIGATIONS	19
3 RESULTS	22
3.1 OVERVIEW OF SCHEMES	22
3.1.1 EPIDERM	22
3.1.2 SWORD	22
3.1.3 THOR-GP	23
3.2 TIME TRENDS BY DISEASE CATEGORY	24
3.2.1 TOTAL WORK-RELATED ILLNESS	24
3.2.2 WORK-RELATED SKIN DISEASE	27
3.2.3 WORK-RELATED RESPIRATORY DISEASE	47
3.2.4 WORK-RELATED MUSCULOSKELETAL DISORDERS	65
3.2.5 WORK-RELATED MENTAL ILL-HEALTH	74
4 DISCUSSION	81
5 CONCLUSION	87

LIST OF TABLES

	Page number
1 Data period for trends analyses	13
2 Categories of illness included in the analyses	15
3 Summary of model features	18
4 Influence of excess zeros on the average annual percentage change in reported incidence in work-related illness	21
5 Average annual percentage change in reported incidence in total work-related illness	25
6 Relative risk by year, with 95% comparison intervals, total work-related illness (core analyses 2009 estimate =1, sample analyses 2014 estimate = 1)	25
7 Average annual percentage change in reported incidence in work-related skin disease	29
8 Relative risk by year, with 95% comparison intervals, total skin disease (EPIDERM 2014 estimate =1, THOR-GP core 2009 estimate = 1, THOR-GP sample 2014 estimate = 1)	30
9 Relative risk by year, with 95% comparison intervals, all contact dermatitis (2014 estimate = 1)	33
10 Relative risk by year, with 95% comparison intervals, allergic contact dermatitis (2014 estimate = 1)	35
11 Relative risk by year, with 95% comparison intervals, irritant contact dermatitis (2014 estimate = 1)	37
12 Relative risk by year, with 95% comparison intervals, mixed contact dermatitis (2014 estimate = 1)	39
13 Relative risk by year, with 95% comparison intervals, contact urticaria (2014 estimate = 1)	41
14 Relative risk by year, with 95% comparison intervals, neoplasia (2014 estimate = 1)	43
15 Relative risk by year, with 95% comparison intervals, other (than contact dermatitis) skin (2014 estimate = 1)	45
16 Average annual percentage change in reported incidence in work-related respiratory disease	48

17	Relative risk by year, with 95% comparison intervals, total respiratory disease (2014 estimate = 1)	49
18	Relative risk by year, with 95% comparison intervals, asthma (2014 estimate = 1)	51
19	Relative risk by year, with 95% comparison intervals, mesothelioma (2014 estimate = 1)	53
20	Relative risk by year, with 95% comparison intervals, benign pleural plaques (2014 estimate = 1)	55
21	Relative risk by year, with 95% comparison intervals, benign pleural plaques – predominantly plaques (2014 estimate = 1)	57
22	Relative risk by year, with 95% comparison intervals, benign pleural plaques – predominantly diffuse (2014 estimate = 1)	59
23	Relative risk by year, with 95% comparison intervals, pneumoconiosis (2014 estimate = 1)	61
24	Relative risk by year, with 95% comparison intervals, other (than those investigated separately) respiratory disease (2014 estimate = 1)	63
25	Average annual percentage change in reported incidence in total work-related musculoskeletal disorders	66
26	Relative risk by year, with 95% comparison intervals, total work-related musculoskeletal disorders (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1)	66
27	Relative risk by year, with 95% comparison intervals, upper limb disorders (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1)	68
28	Relative risk by year, with 95% comparison intervals, spine/back disorders (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1)	70
29	Relative risk by year, with 95% comparison intervals, lower limb disorders (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1)	72
30	Average annual percentage change in reported incidence in total work-related mental ill-health	74
31	Relative risk by year, with 95% comparison intervals, total mental ill-health (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1)	75

32	Relative risk by year, with 95% comparison intervals, anxiety and depression (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1)	77
33	Relative risk by year, with 95% comparison intervals, other work stress (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1)	79

LIST OF FIGURES

	Page number
1 Relative risk by year (core analyses 2009 estimate =1, sample analyses 2014 estimate = 1), with 95% comparison intervals, total work-related illness	26
2 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, total skin	31
3 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, all contact dermatitis	34
4 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, allergic contact dermatitis	36
5 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, irritant contact dermatitis	38
6 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, mixed contact dermatitis	40
7 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, contact urticaria	42
8 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, neoplasia	44
9 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, other (than contact dermatitis) skin	46
10 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, total respiratory disease	50
11 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, asthma	52
12 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, mesothelioma	54

13	Relative risk by year (2014 estimate = 1), with 95% comparison intervals, benign pleural plaques	56
14	Relative risk by year (2014 estimate = 1), with 95% comparison intervals, benign pleural plaques – predominantly plaques	58
15	Relative risk by year (2014 estimate = 1), with 95% comparison intervals, benign pleural plaques – predominantly diffuse	60
16	Relative risk by year (2014 estimate = 1), with 95% comparison intervals, pneumoconiosis	62
17	Relative risk by year (2014 estimate = 1), with 95% comparison intervals, other (than those investigated separately) respiratory disease	64
18	Relative risk by year (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1), with 95% comparison intervals, total musculoskeletal disorders	67
19	Relative risk by year (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1), with 95% comparison intervals, upper limb disorders	69
20	Relative risk by year (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1), with 95% comparison intervals, spine/back disorders	71
21	Relative risk by year (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1), with 95% comparison intervals, lower limb disorders	73
22	Relative risk by year (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1), with 95% comparison intervals, total mental ill-health	76
23	Relative risk by year (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1), with 95% comparison intervals, anxiety and depression	78
24	Relative risk by year (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1), with 95% comparison intervals, other work related stress	80

LIST OF APPENDICES

	Page number
A Summary of reporter fatigue investigations	91
B Descriptive analyses	94

1. BACKGROUND

An objective of occupational disease surveillance is to be able to assess change in incidence of work-related illness (WRI) over time. One approach is to simply look at case counts over time but this method does not take into account other factors that might influence the trend (for example, changes in the numbers of reporters). Thus, in 2005, McNamee *et al*¹ proposed a methodology (in a report submitted to the UK Health and Safety Executive (HSE)) to assess change in incidence of WRI over time using surveillance data collected by The Health and Occupation Research (THOR) network². This method proposed using a multi-level model (MLM) which enables change over time in the number of reporters and in other reporter characteristics which could independently impact on case density to be taken into account. This method was subsequently employed to determine trends in incidence for the period 1996 to 2004³, and in agreement with HSE, on an annual basis thereafter, thus incorporating each additional year of available data⁴⁻¹¹.

Extensive work in trends analyses in THOR data has successfully addressed in turn a range of methodological issues. Most recently the issue of 'reporter fatigue' which has been the main residual methodological issue within this project is being addressed. During this period, additional analyses have been undertaken (and reported on) investigating whether physicians participating in THOR are exhibiting 'reporter fatigue', and if so, how it impacts on the estimate of trend and whether it can be adjusted for^{3, 6, 12-15}. The culmination of this body of work (to date) is currently being written up for peer review¹⁶ and it has been agreed with HSE that the annual trends estimates will not be formally adjusted (if appropriate) for fatigue until after this process. However, the importance of the implications of this body of work regarding the interpretation of the annual trends estimates is also recognised. To this end, the present report includes a summary of the work in this area, including guidance on interpreting the annual trends results if the existing evidence for fatigue from this work was accepted.

This latest report, therefore, describes the trend in incidence of WRI based on data from the three THOR schemes currently supported by the Health and Safety Executive (HSE): case reports of work-related skin disease reported to EPIDERM by dermatologists (1996-2014), case reports of work-related respiratory disease reported to SWORD by chest physicians (1999-2014), and case reports of (any) WRI reported to THOR-GP by general practitioners (2006-2014).

2. METHOD

A full description of the methodology employed in this study is provided hereunder.

2.1 DATA PERIOD

The data period used for the trends analysis is shown in Table 1.

Table 1 Data period for trends analyses

	Scheme start date	Data period for trends study		
		All reporters	Core reporters	Sample reporters
EPIDERM	1993	1996-2014	1996-2014	1996-2014
SWORD	1989	1999-2014	1999-2014	1999-2014
THOR-GP	June 2005	N/A	2006-2009	2011-2014

2.2 REPORTER GROUPS

Physicians reporting to THOR report either as core reporters (reporting every month) or as sample reporters (reporters who report one randomly allocated month a year). The composition of each of the schemes is as follows:

EPIDERM: Consultant dermatologists began reporting to EPIDERM in 1993 and initially all reporters reported at 3-month intervals¹⁷. In January 1996 the scheme was redesigned to consist of a core group with a special interest in occupational skin disease who reported to the scheme on a monthly basis (24 dermatologists originally) with the remaining specialists (220 originally) assigned to report on a sample basis. This mix of core and sample reporters i.e. a smaller core group consisting generally of 'keen specialists' and a larger sample group, continued for the period covered by the current report (1996-2014). For this scheme, analyses based on all reporters combined and separately for core and sample groups were carried out.

SWORD: UK wide SWORD reporting began in 1989¹⁸ and originally physicians could report either monthly (78% of physicians originally), quarterly (19%), bi-annually (<1%) or annually (2%). This original system of reporting was modified in January 1992 (to combat potential reporter fatigue) with those physicians who had reported the most cases forming a core group (approximately 10% of physicians at that time) with the remainder assigned to report on a sample (monthly) basis. As for EPIDERM, this structure of a smaller group of keen specialists and a larger sample group continued throughout the time period covered by these analyses (1999-2014 for SWORD). For this scheme, analyses based on all reporters combined and separately for core and sample groups were carried out.

THOR-GP: Initially all reporters to THOR-GP reported on a core basis. This permitted the scheme to 'come up to speed' rapidly, and to provide the HSE with

early data on the distribution of GP reported WRI from THOR-GP. Sample reporting was introduced to THOR-GP in April 2007 to counteract possible reporter fatigue with a small number of sample reporters (33 (7%) reporters during the period April 2007 to December 2009). In 2010, the proportion of sample reporters increased to 78% to give a core:sample ratio of 1:4. Since the introduction of sample reporting in 2007, all new reporters that were recruited to the scheme were randomly allocated to participate as either core or sample. Additionally, in 2010 all reporters not previously assigned core or sample status were randomly allocated to one of these two different reporting groups. From 2011 onwards, (especially because HSE could no longer fund remuneration for GP participation) all reporters became sample reporters. Because of these extensive changes (there is evidence that GPs might report more cases during any one month compared to core reporters)¹⁹, it was not felt appropriate to consider THOR-GP trends for the period as a whole, for all reporters combined (core and sample) or to directly compare pre and post 2010/2011 trends. Therefore GP analyses were carried out for sample reporters only for the period 2011 onwards, and for core reporting only for the period 2006-2009. Given that core reporting has now ceased, it is the former that will be most informative of future trends for this group.

Definition of an active reporter: For the purpose of the analyses it was deemed important to include only those reporters with evidence of active participation. For the THOR specialist schemes an active reporter was defined as a reporter who either returned cases or declared 'I have nothing to report' (a zero return) during the study period. For THOR-GP, reporters can submit a sickness absence (SA) return only in any given month (i.e. information about additional sickness absence that has been issued to a previously reported case). Approximately 5 reporters a month submit a SA return with no other information about cases (case or zero return): ideally they should also have submitted a zero return if there are no new cases. While these reporters are, in the general sense of the word, active, in terms of contributing information about incidence they are not. On the other hand, it might be assumed that, if they had seen new cases they would have contributed them and therefore this corresponds to a zero return in terms of incidence. However, as it is difficult to be sure of this and this activity accounts for a very small proportion of the monthly returns, we considered them to be inactive. Therefore for the purpose of this trends analysis also, a THOR-GP reporter has to have submitted a case or zero return to be considered active for the purpose of studying trends in incidence.

2.3 CATEGORIES OF ILLNESS

Initial power calculations undertaken for the THOR specialist schemes suggested that a specific disease category should only be investigated (separately) if the number of actual cases reported during the study period exceeded 250³. For THOR-GP it was decided that, although over a shorter time period, the minimum number of

cases required for any disease category to be included in the analysis would remain at 250. The resulting disease groups to be included in the analysis are shown in Table 2.

Table 2 Categories of illness included in the analyses

	Clinical specialist	THOR-GP
All work-related illness	-	Yes
Total skin	Yes	Yes
Contact dermatitis (CD)	Yes	-
• Allergic CD	Yes	-
• Irritant CD	Yes	-
• Mixed CD	Yes	-
Neoplasia	Yes	-
Contact urticaria	Yes	-
Other skin (other than contact dermatitis)	Yes	-
Total respiratory	Yes	-
Asthma	Yes	-
Mesothelioma	Yes	-
Benign pleural disease	Yes	-
• Predominantly plaques	Yes	
• Predominantly diffuse	Yes	
Pneumoconiosis	Yes	-
Other respiratory disease (other than those specified above)	Yes	-
Total musculoskeletal	-	Yes
Upper limb disorders (hand/wrist/arm/shoulder/elbow)	-	Yes
Spine/back disorders (neck/thoracic spine/lumbar spine/trunk)	-	Yes
Lower limb disorders (hip/knee/ankle/foot)	-	Yes
Total mental ill-health	-	Yes
Anxiety and depression	-	Yes
Other work stress	-	Yes

2.4 THE MULTI-LEVEL MODEL AND ITS ASSUMPTIONS

The STATA software command **xtnbreg** was used to fit longitudinal, negative binomial (i.e. over-dispersed) Poisson models with random effects.

In these models, the dependent variable was the number of actual cases, including zeros, per reporter per month; the main 'covariate' is calendar time. The aim of the analysis is to estimate the relationship between annual UK incidence rate and time, after adjusting for potential confounders. Numbers of cases might vary from year to year solely because of changes in the size of the UK working population, even though the rate is constant. Therefore estimated population sizes for each year (see below) were included in the model as an 'offset'; this feature means that the model estimates change in rates, not changes in case counts.

Apart from 'calendar time', the other variables included in the regression models as covariates were 'season', 'reporter type' (core or sample), 'first month/s as a new reporter'. These are factors that can influence the reported incidence levels. Further details of covariates/offsets in the model are given later in this section.

It is important to allow for the possible impact of having different reporting centres at different periods of time: some centres may have a larger, or more 'at risk' catchment patient population than others. In a statistical model, we can take account of such differences by allowing the incidence level to vary between centres; the analysis can then trace the pattern over time 'within centres'. In a 'fixed effects' approach to this, the incidence level is estimated for each centre; in a 'random effects' model, the incidence levels are assumed to vary randomly between centres in each subgroup (e.g. subgroups of core reporters and sample reporters) but not estimated directly. In previous reports, two sets of results were presented corresponding to each of these options but, after consultation with HSE, it was decided that from 2010 onwards only results based on models with random effects would be presented. (One reason was because the fixed effects model omits all reporters who had reported only zero cases throughout the study period).

Every statistical model has to make an assumption about the form of the variability which remains after taking into account all covariates in the model. The Poisson distribution is the usual distribution assumed for count data; the Negative Binomial distribution is a more general version of a Poisson distribution which is less rigid; in the Poisson the variance and mean are constrained to be equal, but not in the Negative Binomial.

Calendar time – For the main analyses, changes in incidence were estimated in two different ways: 1) '*non-parametric*' approach: the model contained separate indicator variables for different years. In the current analyses, 2014 was taken as the reference year (2009 for THOR-GP core only analyses) and the percentage increase or decrease in incidence compared to 2014 (or 2009) was estimated. These analyses had no in-built assumptions about the pattern of change over time. 2) '*parametric*' approach with a continuous time variable measured on a scale of years. The statistical models for these analyses assumed a systematic trend throughout the period being studied. Specifically, it was assumed that the percentage change from

one year to the next is a constant throughout the relevant period. Where the assumption is valid, this parametric approach offers a more precise way of estimating change than approach 1.

Season – Seasonal variation refers to variation within a year whose pattern tends to be repeated from year to year. This short-term variation could be due to seasonal variation in illness or seasonality in reporting behaviour; the latter could occur because of holidays, for example. To address this, indicator variables for months (with June as the reference category) were included in the models. Seasonal variation should not bias the assessment of long-term changes in this study. However it could affect precision in the estimate of trend if not controlled.

Reporter type – Reporter type (core or sample) had been shown to cause variation in incidence between reporters. Thus, a variable which took the value '1' if a core reporter and '0' if a sample reporter was included in the models. Furthermore, for the purpose of the analysis, if a reporter changed from the core reporting group to sample reporting or vice versa, he or she was treated as a new reporter for the period after the change. We have previously shown¹⁹ that there are differences in behaviour for the same reporter depending on whether they are reporting as core or as sample.

First month/s as a new reporter – It is conceivable that, in the first month/s of reporting, a new entrant to a surveillance scheme might include cases seen over a period longer than the assigned single month. If there was a sufficiently large 'harvest' of old cases, it could produce a false, decreasing 'trend' over time. For the THOR specialist schemes, initial investigations suggested that 'new recruit' harvesting might be occurring during the first month that a reporter actively reported to a scheme. Thus, to control for harvesting, a variable which took the value '1' if it was the first month the reporter had reported and '0' for all other months was included in the models. Initial investigations suggested the period of 'harvesting' maybe longer for THOR-GP compared to the specialist schemes (5 months compared to 1). This might occur because, compared to specialists, there is more opportunity for 'old' cases to present themselves again to a GP, thus prompting a report. Thus, variables representing the first 5 months of active reporting were included in the THOR-GP regression models. Moreover, for these main analyses, the first 7 months of THOR-GP were excluded (June to December 2005). Since approximately 28% of the GPs (reporting between 2006 and 2013) joined the scheme in 2005, it was felt that the 2005 data may have been particularly prone to the effect of harvesting. Of note, for those reporters joining THOR-GP as sample reporters (19% of the total ever reporters), the period of harvesting was taken as 1 month rather than 5 (as the latter would in effect, equate to 5 years).

Population change - Analysis of data from the UK Labour Force Survey (LFS) had shown a fairly regular increase in the size of the working population of the order of

1% a year up to 2006²⁰, although decreases may have occurred since then. One might perhaps expect to see an increase in cases over time because of this even if true incidence *rates* remained constant. Therefore we have accounted for this change in population base by including in the ML model an offset variable representing the UK working population, obtained from the LFS, for each year.

Table 3 Summary of model features

Feature	Description
Centre variation	Variation in incidence between centres is assumed; analysis attempts to measure change within centres
Centre number	If a reporter changed from core to sample (or vice versa) they were assigned a new centre number and thus treated as a new reporter in the model
Denominators/population sizes	The catchment population for each centre is assumed to increase/decrease in line with changes in the size of UK working population
Unexplained variation	Assumed to follow a Negative Binomial distribution
Active reporter	Only 'active' reporters were included in the analysis. This was defined as a reporter who either returned cases or declared 'I have nothing to report' (a zero return) at least once during the study period.
New recruit 'harvesting' of old cases	For SWORD and EPIDERM, the model assumes that this effect only occurs during the first month of reporting or the first month a reporter returned as a core reporter. For THOR-GP, it allows it to occur for the first 5 months of reporting for those joining as core and 1 month for those joining as sample reporters.
Calendar time treatment: non-parametric approach	Rate Ratio for each year compared to 2013 is estimated
Calendar time treatment: parametric approach	A linear trend over time is assumed: Rate Ratio for each year compared to the previous one is estimated

2.5 SUMMARY OF 'REPORTER FATIGUE' INVESTIGATIONS

As membership time increases, a reporter might become less committed to active participation but still retain membership. How such 'fatigue' manifests itself and whether this can cause bias in time trend estimation has been a major methodological concern for this project. Previous investigations reported to HSE have focussed on two different manifestations of fatigue; an increase in non-response over time and an increase in zero (blank) returns over time^{3, 6, 12-15}. We have argued previously that an increase in non-response over time would not necessarily cause bias in trends estimation; therefore results of these analyses have not been reproduced here. In contrast, an increase in zero returns over time, some of which may be 'false zeroes' and which do not truly equate to 'zero cases', would mean that the trend over time would be biased downwards compared to the situation if there were no reporting fatigue.

Steps taken to investigate this particular manifestation of fatigue are summarised in Appendix 1. The most recent (and we believe improved) approach has been the application of a zero-inflated negative binomial (ZINB) model. These analyses have recently been written up in an article submitted for peer review¹⁶ with an overview provided below.

Analysis of zero-inflated count data using a zero-inflated negative binomial model (ZINB)

To account for the presence of excess zero cases within the reported data, the reported monthly number of cases was fitted using a Zero-Inflated Negative Binomial Model (ZINB) with multi-level random effects.

This model has two parts; the first supposes that, on occasion, a reporter might send back a zero report regardless of the actual number of cases seen i.e. an excess zero. This part of the model supposes a binary decision: send back an excess zero regardless or send back the true count zero or otherwise. The second part is the usual negative binomial model for true cases, including true zero cases, each month. The model allows for two sets of predictors in the two portions of the model. These were mean centred membership year (first part of model) and calendar time (second part of model). Thus the complete model allows for the possibility of excess zeros in the data; it can estimate their frequency and can estimate the true trend after allowing for this phenomenon.

The covariate thought to influence zero case reports and therefore included in the first part of the model was peak holiday season. Covariates thought to influence the incidence of work-related illness, and therefore included in the second part of the model, were first month as a reporter and months of the year containing a bank

holiday. All modelling was repeated for 'core' reporters only, 'sample' reporters only, and both 'core' and 'sample' reporters.

Using this approach, data for EPIDERM (1996-2012), SWORD (1999-2012) and THOR-GP (2006-2012) were analysed¹⁴⁻¹⁶. The impact of adjusting for excess zeros on the annual average percentage change in incidence of total work-related skin disease (EPIDERM), total work-related respiratory disease (SWORD) and total WRI (THOR-GP) is shown in Table 4.

Results:

EPIDERM

The results suggest that both core and sample dermatologists reporting to EPIDERM are exhibiting reporter fatigue. Overall core reporters were less likely to report an excess zero than sample, yet both experienced an increase in excess zero returns with increasing membership time. Thus, adjusting for 'excess zeros' would have a greater impact on the trend estimates for sample reporters compared to core. However, because sample reporters contribute less data, the impact on the overall estimate (core and sample) is less pronounced.

SWORD There is little evidence that SWORD core reporters are exhibiting reporter fatigue as shown by an increase in excess zero returns with increasing membership time. The evidence of reporting fatigue for SWORD sample reporters appears to be less strong than for EPIDERM sample reporters but there does appear to be fatigue manifesting in this way for this group. For SWORD, sample reporters contribute more data than core reporters and therefore fatigue in this group may have more impact on the overall estimate (compared to core).

THOR-GP There was some evidence of reporter fatigue (as shown by an increase in excess zero returns with increasing membership time) for THOR-GP. Contrary to SWORD and EPIDERM, excess zeros were more likely to be reported by core compared to sample reporters. Since core reporting has now ceased, it is the impact of fatigue on sample reporting that is of greater interest. It should be noted that sample reporting is a relatively new phenomenon in THOR-GP (only 100% since 2011) and therefore it is important to continue to monitor the potential impact of reporter fatigue on this group.

Table 4 **Influence of excess zeros on the average annual percentage change in reported incidence in work-related illness**

		Core	Sample	Core + sample
EPIDERM	Member year ^a	1.14 (1.06, 1.22)*	1.09 (1.05, 1.12)*	1.08 (1.05, 1.12)*
(Total skin disease)	Negative binomial ^b	-2.8	-1.8	-2.6
	ZINB ^c	-2.4	0.0	-2.3
	% change ^d	14%	100%	12%
	Vuong p-value ^e	<0.001	0.003	<0.001
SWORD	Member year	1.04 (0.94, 1.14)	1.05 (1.02, 1.08)*	1.04 (1.02, 1.07)*
(Total respiratory disease)	Negative binomial	-2.7	-2.4	-2.5
	ZINB	-2.8	-0.5	-2.1
	% change	4%	79%	16%
	Vuong p-value	0.406	0.053	0.012
THOR-GP	Member year	1.26 (1.11, 1.44)*	1.10 (0.92, 1.32)	1.17 (1.05, 1.29)*
(Total work-related illness)	Negative binomial	-13.8%	-23.7%	-14.4%
	ZINB	-9.2%	-20.9%	-10.98%
	% change	34%	12%	25%
	Vuong p-value	0.010	0.263	0.006

*Statistically significant at the 5% level or below

^aExcess zero odds ratio: This denotes whether the proportion of excess zeros is (significantly) increasing with membership time. For example, for EPIDERM core reporters, excess zeros increase by 14% per year of membership and this increase is statistically significant

^bAnnual average percentage change in incidence from negative binomial model (i.e. not adjusted for excess zeros)

^cAnnual average percentage change in incidence from zero-inflated negative binomial model (i.e. adjusted for excess zeros)

^dPercentage difference between negative binomial model and zero-inflated negative binomial model

^eVuong test comparing whether the zero-inflated negative binomial model is a statistically better fit to the data than the negative binomial model

3 RESULTS

3.1 OVERVIEW OF SCHEMES

An overview of the reporting activity of the physicians participating in EPIDERM, SWORD and THOR-GP is provided in Appendix B and briefly described below.

3.1.1 EPIDERM

A total of 449 dermatologists have been enrolled in EPIDERM during the study period with 93% actively participating at least once (i.e. either returning cases or declaring 'I have nothing to report this month'). On average, 193 dermatologists participated in EPIDERM each year and 2014 saw a small drop in the overall number of physicians in EPIDERM (from 165 in 2013 to 153 in 2014). Response rates (cards returned/cards sent out) per year showed an initial increase between 1996 and 2001, after which they exhibited an overall decline until 2011 when they stabilised at around 60%. The number of active reporters per month has shown a similar pattern with an average of 22 per month in 2014. The average cases per active reporter increased slightly between 2013 and 2014 (1.9 in 2013 and 2.0 in 2014). Reporters to EPIDERM are predominantly sample (86% in 2014) but core reporters report more cases per active reporter per month (3.5) compared to sample (0.9). Case reports to EPIDERM continue to be predominantly contact dermatitis (82% of total cases) with smaller proportions of neoplasia (12%) and other skin diagnoses.

3.1.2 SWORD

Active participation in SWORD during the study period was similar to EPIDERM with 94% of the 861 chest physicians enrolled during this period actively reporting at least once. On average, 471 chest physicians participated in SWORD each year and the total number of reporters in SWORD decreased slightly between 2013 and 2014 (447 to 433). Response rates (cards returned/cards sent out) showed a decrease between 1999 and 2006 but have remained relatively constant (60-70%) thereafter. The average number of active reporters per month and the average number of cases per active reporter decreased slightly between 2013 and 2014 (2013: 32 and 1.2; 2014: 29 and 1.1, respectively). Similar to EPIDERM, the smaller group of chest physicians reporting as core reported more cases per active reporter per month than chest physicians reporting as sample. The majority of the diagnoses (43%) reported to SWORD during the study period were benign pleural plaques. Of the remaining cases 19% were mesothelioma, 19% asthma, 9% pneumoconiosis, and 14% 'other' respiratory disease.

3.1.3 THOR-GP

THOR-GP differs from EPIDERM and SWORD in that at the start of the study period (2006) all GPs reported to THOR-GP on a core basis whilst since 2011, reporting has been exclusively sample, with the majority of the switch from core to sample occurring in 2010. On average, 263 GPs were enrolled in THOR-GP each year. Active participation during this period was lower than observed for SWORD and EPIDERM (77% of the 549 GPs ever enrolled). In contrast to the clinical specialists, GPs participating as sample physicians reported on average more cases per active reporter per month than GPs participating as core reporters. The average number of active reporters per month has decreased slightly since the introduction of 100% sample reporting (15 in both 2011 and 2012, 14 in 2013, 13 in 2014), as has the average number of cases per active reporter month (1.4 for 2011 and 2012, 1.3 for 2013, 1.1 in 2014). Musculoskeletal and mental ill-health case reports comprised the majority (52% and 32%, respectively) of the cases reported to THOR-GP with smaller proportions of skin (10%), respiratory (3%) and other diagnoses (6%).

3.2 TIME TRENDS BY DISEASE CATEGORY

This report continues with the approach first adopted in the trends report submitted to HSE in September 2010⁷, in that the statistical uncertainty (confidence intervals) in the graphs illustrating time trends are presented in such a way as to allow the reader to assess the significance of the difference between any two years. This useful approach suggested by the then HSE liaison officer (John Hodgson) when steering the research follows the method described by Firth and de Menezes²¹ which assigns a confidence (or comparison) interval to the reference category (2014 in the present analyses) and reduces the width of the confidence (comparison) intervals of non-reference categories in such a way that all pairwise comparisons between years can validly be made using these adjusted confidence intervals.

Interpreting the results in light of reporter fatigue

The results presented here have not been formally adjusted for reporter fatigue. However, for SWORD and EPIDERM the likely impact of adjusting for 'excess zeros' (i.e. if a zero inflated binomial model was applied to the data) has been *estimated* by applying the results of the analyses described under Section 2.5 (Table 4) to the present data. These analyses suggested that adjusting for excess zeros would reduce the annual average estimated decrease from -2.6% to -2.3% for EPIDERM and from -2.5% to -2.1% for SWORD.

Thus, for example if an annual average decrease in incidence of -3.8% was observed in the present study for total skin disease reported to EPIDERM, adjusting this for the likely impact of fatigue based on the above would reduce the decrease to approximately 3.4%. These adjustments have been carried out for total skin (EPIDERM), total respiratory (SWORD) and total WRI (THOR-GP) only (it cannot be assumed at this stage that the observed effect would be the same across the different diagnoses).

Note: it must be stressed that these are estimates, provided for guidance purposes only.

3.1.4 TOTAL WORK-RELATED ILLNESS

The average annual percentage change in risk of total WRI, as reported to THOR-GP is shown in Table 5 whilst the relative rates by year are shown in Tables 6 and Figure 1. Based on reports from core reporters only, the average annual decrease in incidence for the period 2006-2009 was -13.5% (95% CIs: -16.2, -10.8). This compares to an annual average decrease of -14.6% (95% CIs: -21.2, -7.5) for analyses based on sample reporters only for the period 2011-2014. The graphs

showing relative risk by year for THOR-GP suggest a year on year decline for both core (2006-2009) and sample reporters (2011-2014). For sample reporters, the data suggest the largest drop in incidence to be occurring between 2013 and 2014.

For estimates based on core reports (2006-2009), extrapolating the results of the previously described analyses using a zero-inflated binomial model (Section 2.5), the impact of adjusting for 'excess zeros' in the present analyses *might be* expected to change the average annual percentage change in incidence of total WRI in the order of from -13.5% to -8.9%. There was no evidence of fatigue manifesting in this way for sample reporters, and as previously stated, THOR-GP is now exclusively reliant on sample reporters (since HSE stopped funding honoraria for participation).

Table 5 Average annual percentage change in reported incidence in total work-related illness

	ESTIMATED % CHANGE (95% CONFIDENCE INTERVAL)	
	THOR-GP	
	Core reporters	Sample reporters
Year (continuous)		
2006-2009	-13.5 (-16.2, -10.8)	/
2011-2014	/	-14.6% (-21.2, -7.5)

Models adjusted for season and harvesting

Population offset included in the model

The number of actual cases on which each analysis is based is provided in Table B6 on page 117

Table 6 Relative risk by year, with 95% comparison intervals, total work-related illness (core analyses 2009 estimate =1, sample analyses 2014 estimate = 1)

	Relative rates (95% comparison interval)	
	THOR-GP	
	Core reporters	Sample reporters
YEAR		
2006	1.68 (1.57,1.79)	/
2007	1.33 (1.25,1.4)	/
2008	1.23 (1.16,1.31)	/
2009	1 (0.93,1.08)	/
2010	/	/
2011	/	1.54 (1.34,1.78)
2012	/	1.42 (1.23,1.63)
2013	/	1.28 (1.1,1.48)
2014	/	1 (0.85,1.18)


Models adjusted for season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B6 on page 117

Figure 1 Relative risk by year (core analyses 2009 estimate = 1, sample analyses 2014 estimate = 1), with 95% comparison intervals, total work-related illness

a) Core reporters



b) Sample reporters



3.1.5 WORK-RELATED SKIN DISEASE

The average annual percentage change in risk of work-related skin disease, as reported by dermatologists and GPs is shown in Table 7 whilst the relative rates by year are shown in Tables 8 to 15 and Figures 2 to 9.

Dermatologists reporting to EPIDERM: The annual average change in incidence of dermatologist reported work-related skin disease (1996-2014) was -3.8% (95% CIs: -4.3, -3.3). This compares to the previous estimate of -3.6% (95% CIs: -4.1, -3.0) reported in 2014 (based on data for the period 1996-2013). The graphs showing relative risk by year suggest an initial decrease in incidence in the earlier part of the study period (1996-2005) followed by a relatively flat trend (2006-2012) and a further decrease between 2012 and 2013. There was little suggestion of a further decrease between 2013 and 2014. The estimated annual change in incidence of contact dermatitis (CD) was similar at -3.7% (95% CIs: -4.3, -3.2) with a similar annual pattern. Analyses of shorter-term trends (2006-2013) for CD suggested an annual average decrease in CD incidence of -3.9% (95% CIs: -5.5, -2.2) per year. Analysis by type of CD indicated a steeper decrease in the incidence of allergic CD (-5.4% (95% CIs: -6.2, -4.7)) compared to irritant CD (-2.5 (95% CIs: -3.2, -1.7)) or mixed CD (-2.5% (95% CIs: -3.7, -1.4)) and these estimates remain relatively unchanged by the addition of the 2014 data. The graphs showing relative risk by year suggest an overall downward trend for allergic CD between 1996 and 2006 followed by a relatively flat trend. For irritant CD, after an initial decrease between 1999 and 2000, the trend is flat until 2012, after which it decreases steeply, with little change between 2013 and 2014. The annual average change in incidence of dermatologist reported urticaria (1996-2013) was -7.1% (95% CIs: -8.8, -5.4) compared to the previously reported -6.6% (95% CIs: -8.4, -4.9) (based on data for 1996-2013). The trend in incidence for neoplasia also suggested a decrease, of -3.0% (95% CIs: -4.5, -1.6) compared to -3.2% (95% CIs: -4.6, -1.7) reported previously.

There was some variation by reporter type (core versus sample). This was most pronounced for neoplasia with data from core reporters suggesting an annual average decrease of -4.6% (95% CIs: -6.2, -3.0) whilst data from sample reporters suggested an increase of 2.6% (95% CIs: -0.5, 5.7) and for urticaria (core: -7.7% (95% CIs: -9.4, -5.9); sample: -0.9 (95% CIs: -7.1, 5.7)).

Extrapolating the results of the previously described analyses using a zero-inflated binomial model (Section 2.5), the impact of adjusting for 'excess zeros' in the present analyses *might be* expected to change the average annual percentage change in incidence of total work-related skin disease in the order of from -3.8% to -3.4%.

GPs reporting to THOR-GP: For GPs the estimated annual decrease in incidence of total work-related skin disease based on reports from core reporters only (2006-2009) was -6.4% (95% CIs: -14.8, -3). For sample reporters only (2011-2014), the estimated annual decrease was -19.2% (95% CIs: -37.8, 4.9). The graph showing relative rates by year suggests a general decrease over time for core reporters (although confidence intervals are overlapping for all years). For sample reporters, there is an initial decrease between 2011 and 2012 followed by a relatively flat trend, although confidence intervals are again very wide and overlapping for all years, since the overall numbers reported by monthly 'sample' reporting only are considerably less than those which were collected through 'core' reporting at the inception of THOR-GP.

Table 7 Average annual percentage change in reported incidence in work-related skin disease

		ESTIMATED % CHANGE (95% CONFIDENCE INTERVAL)				
		All reporters	Core reporters		Sample reporters	
		EPIDERM	EPIDERM	THOR-GP	EPIDERM	THOR-GP
	Year (continuous)					
Total skin	1996-2014	-3.8 (-4.3, -3.3)	-3.9 (-4.4, -3.4)	/	-2.6 (-4.1, -1.1)	/
	2006-2009	/	1.5 (-2.4, 5.7)	-6.4 (-14.8, -3)	/	/
	2011-2014	/	/	/	-7.2 (-20.4, 8.3)	-19.2 (-37.8, 4.9)
Contact dermatitis (CD)	1996-2014	-3.7 (-4.3, -3.2)	-3.7 (-4.3, -3.2)	/	-3.8 (-5.5, -2.2)	/
	2006-2014	-3.9 (-5.5, -2.2)	/	/	/	/
• Allergic CD	1996-2014	-5.4 (-6.2, -4.7)	-5.7 (-6.5, -4.9)	/	-3.4 (-5.6, -1.1)	/
• Irritant CD	1996-2014	-2.5 (-3.2, -1.7)	-2.4 (-3.2, -1.6)	/	-2.8 (-5.2, -0.4)	/
• Mixed CD	1996-2014	-2.5 (-3.7, -1.4)	-2.8 (-3.9, -1.6)	/	-1.0 (-4.7, 2.8)	/
Urticaria	1996-2014	-7.1 (-8.8, -5.4)	-7.7 (-9.4, -5.9)	/	-0.9 (-7.1, 5.7)	/
Neoplasia	1996-2014	-3.0 (-4.5, -1.6)	-4.6 (-6.2, -3.0)	/	2.6 (-0.5, 5.7)	/
Other* skin	1996-2014	-5.5 (-6.5, -4.5)	-6.6 (-7.6, -5.5)	/	-0.2 (-2.6, 2.3)	/

*Other than contact dermatitis

Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model

The number of actual cases on which each analysis is based is provided in Table B2 on page 99

Table 8 Relative risk by year, with 95% comparison intervals, total skin disease (EPIDERM 2014 estimate = 1, THOR-GP core 2009 estimate = 1, THOR-GP sample 2014 estimate = 1)

	Relative risk (95% comparison interval)				
	All reporters	Core reporters		Sample reporters	
	EPIDERM	EPIDERM	THOR-GP	EPIDERM	THOR-GP
YEAR					
1996	2.14 (1.97,2.32)	2.27 (2.08,2.48)	/	1.32 (0.97,1.79)	/
1997	2.27 (2.11,2.44)	2.37 (2.19,2.56)	/	1.92 (1.48,2.49)	/
1998	2.05 (1.9,2.21)	2.15 (1.98,2.33)	/	1.68 (1.33,2.13)	/
1999	2.09 (1.93,2.25)	2.16 (1.99,2.35)	/	1.85 (1.52,2.27)	/
2000	1.95 (1.81,2.11)	2.08 (1.91,2.26)	/	1.46 (1.17,1.83)	/
2001	1.78 (1.65,1.92)	1.85 (1.71,2.01)	/	1.63 (1.31,2.02)	/
2002	1.74 (1.62,1.88)	1.81 (1.67,1.96)	/	1.6 (1.27,2.01)	/
2003	1.75 (1.62,1.89)	1.87 (1.73,2.03)	/	1.18 (0.91,1.54)	/
2004	1.61 (1.49,1.74)	1.68 (1.55,1.82)	/	1.37 (1.09,1.73)	/
2005	1.61 (1.48,1.74)	1.69 (1.55,1.84)	/	1.28 (1.01,1.63)	/
2006	1.45 (1.33,1.57)	1.47 (1.35,1.61)	1.32 (1.07,1.62)	1.53 (1.24,1.9)	/
2007	1.5 (1.38,1.63)	1.59 (1.45,1.74)	1.21 (1.01,1.45)	1.14 (0.89,1.46)	/
2008	1.4 (1.28,1.53)	1.5 (1.36,1.65)	1.22 (1.02,1.45)	0.98 (0.73,1.31)	/
2009	1.59 (1.45,1.73)	1.66 (1.51,1.82)	1 (0.8,1.24)	1.34 (1.02,1.75)	/
2010	1.49 (1.35,1.64)	1.56 (1.41,1.72)	/	1.26 (0.95,1.68)	/
2011	1.22 (1.09,1.37)	1.24 (1.1,1.41)	/	1.21 (0.89,1.63)	1.66 (1.1,2.51)
2012	1.36 (1.22,1.53)	1.39 (1.23,1.58)	/	1.33 (1.01,1.75)	1.02 (0.6,1.73)
2013	1.04 (0.92,1.18)	1.03 (0.9,1.18)	/	1.15 (0.86,1.54)	0.79 (0.43,1.43)
2014	1 (0.88,1.13)	1 (0.87,1.15)	/	1 (0.73,1.37)	1 (0.59,1.71)


Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B2 on page 99

Figure 2 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, total skin


a) EPIDERM, all reporters




b) EPIDERM, core reporters



c) EPIDERM, sample reporters



d) THOR-GP, core reporters (note scale change)



e) THOR-GP, sample reporters (note scale change)




Table 9 Relative risk by year, with 95% comparison intervals, all contact dermatitis (2014 estimate = 1)


	Relative risk (95% comparison interval)		
	All reporters	Core reporters	Sample reporters
	EPIDERM	EPIDERM	EPIDERM
YEAR			
1996	2.09 (1.91,2.29)	2.17 (1.97,2.4)	1.52 (1.09,2.11)
1997	2.23 (2.05,2.42)	2.31 (2.12,2.51)	2.05 (1.55,2.7)
1998	2.05 (1.89,2.23)	2.13 (1.95,2.33)	1.88 (1.46,2.43)
1999	2.1 (1.93,2.28)	2.12 (1.94,2.32)	2.26 (1.83,2.79)
2000	1.88 (1.72,2.05)	1.96 (1.78,2.15)	1.63 (1.29,2.07)
2001	1.69 (1.55,1.84)	1.77 (1.61,1.93)	1.46 (1.14,1.87)
2002	1.76 (1.63,1.92)	1.82 (1.67,1.99)	1.69 (1.32,2.16)
2003	1.73 (1.59,1.88)	1.84 (1.68,2.01)	1.21 (0.91,1.61)
2004	1.58 (1.45,1.71)	1.66 (1.53,1.81)	1.21 (0.91,1.6)
2005	1.56 (1.43,1.7)	1.62 (1.47,1.78)	1.34 (1.04,1.74)
2006	1.39 (1.27,1.52)	1.44 (1.31,1.59)	1.3 (1,1.68)
2007	1.42 (1.29,1.56)	1.52 (1.37,1.68)	0.93 (0.69,1.25)
2008	1.37 (1.24,1.51)	1.47 (1.32,1.63)	0.89 (0.64,1.24)
2009	1.57 (1.43,1.72)	1.65 (1.5,1.82)	1.16 (0.85,1.59)
2010	1.5 (1.36,1.66)	1.58 (1.42,1.76)	1.15 (0.84,1.59)
2011	1.23 (1.1,1.39)	1.26 (1.11,1.43)	1.21 (0.88,1.68)
2012	1.36 (1.21,1.53)	1.4 (1.24,1.6)	1.25 (0.92,1.7)
2013	1.04 (0.91,1.18)	1.03 (0.9,1.19)	1.12 (0.81,1.55)
2014	1 (0.88,1.14)	1 (0.87,1.15)	1 (0.71,1.4)

Models adjusted for reporter type (where appropriate), season and harvesting
Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B2 on page 99

Figure 3 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, all contact dermatitis

a) EPIDERM, all reporters



b) EPIDERM, core reporters



c) EPIDERM, sample reporters




Table 10 Relative risk by year, with 95% comparison intervals, allergic contact dermatitis (2014 estimate = 1)

	Relative risk (95% comparison interval)		
	EPIDERM		
	All reporters	Core reporters	Sample reporters
YEAR			
1996	2.08 (1.82,2.38)	2.2 (1.92,2.53)	1.47 (0.94,2.3)
1997	2.51 (2.24,2.8)	2.6 (2.32,2.91)	2.19 (1.52,3.17)
1998	2.05 (1.82,2.31)	2.18 (1.93,2.47)	1.4 (0.93,2.09)
1999	2.04 (1.81,2.3)	2.11 (1.86,2.4)	1.84 (1.31,2.59)
2000	2.02 (1.79,2.27)	2.13 (1.88,2.42)	1.57 (1.12,2.2)
2001	1.62 (1.43,1.83)	1.75 (1.54,2)	0.98 (0.65,1.48)
2002	1.84 (1.64,2.06)	1.87 (1.66,2.12)	1.85 (1.33,2.57)
2003	1.8 (1.6,2.02)	1.89 (1.67,2.13)	1.39 (0.96,2.03)
2004	1.42 (1.25,1.61)	1.5 (1.31,1.7)	1.06 (0.69,1.62)
2005	1.43 (1.26,1.64)	1.45 (1.26,1.67)	1.46 (1.02,2.08)
2006	1.35 (1.18,1.54)	1.37 (1.19,1.58)	1.36 (0.95,1.95)
2007	1.16 (1.1,1.35)	1.19 (1.01,1.39)	1.11 (0.74,1.66)
2008	1.19 (1.02,1.38)	1.23 (1.05,1.45)	1.01 (0.64,1.58)
2009	1.16 (0.99,1.35)	1.21 (1.03,1.42)	0.88 (0.53,1.48)
2010	1.15 (0.98,1.35)	1.16 (0.98,1.38)	1.18 (0.75,1.88)
2011	0.89 (0.73,1.08)	0.85 (0.69,1.05)	1.29 (0.8,2.06)
2012	1.09 (0.91,1.31)	1.14 (0.93,1.38)	0.92 (0.55,1.53)
2013	0.98 (0.81,1.19)	0.96 (0.78,1.19)	1.06 (0.66,1.68)
2014	1 (0.83,1.21)	1 (0.81,1.23)	1 (0.61,1.64)


Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B2 on page 99

Figure 4 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, allergic contact dermatitis

a) EPIDERM, all reporters



b) EPIDERM, core reporters



c) EPIDERM, sample reporters




Table 11 Relative risk by year, with 95% comparison intervals, irritant contact dermatitis (2014 estimate = 1)

	Relative risk (95% comparison interval)		
	EPIDERM		
	All reporters	Core reporters	Sample reporters
YEAR			
1996	2.19 (1.93,2.47)	2.38 (2.09,2.72)	0.88 (0.48,1.59)
1997	2.05 (1.83,2.31)	2.24 (1.98,2.53)	1.28 (0.78,2.09)
1998	2.12 (1.89,2.38)	2.25 (1.99,2.54)	1.79 (1.25,2.57)
1999	2.23 (1.98,2.5)	2.38 (2.1,2.7)	1.75 (1.25,2.46)
2000	1.81 (1.59,2.06)	1.96 (1.71,2.25)	1.29 (0.88,1.9)
2001	1.73 (1.53,1.95)	1.81 (1.59,2.07)	1.65 (1.18,2.3)
2002	1.76 (1.56,1.99)	1.9 (1.68,2.16)	1.38 (0.93,2.04)
2003	1.7 (1.5,1.92)	1.86 (1.63,2.11)	1.08 (0.7,1.68)
2004	1.52 (1.34,1.72)	1.66 (1.45,1.89)	1.03 (0.66,1.59)
2005	1.71 (1.51,1.94)	1.85 (1.62,2.12)	1.19 (0.79,1.78)
2006	1.58 (1.4,1.8)	1.74 (1.53,1.99)	1.14 (0.76,1.7)
2007	1.76 (1.56,2)	2 (1.76,2.27)	0.72 (0.45,1.18)
2008	1.61 (1.41,1.85)	1.8 (1.57,2.07)	0.8 (0.47,1.36)
2009	1.81 (1.59,2.05)	2.02 (1.78,2.3)	0.78 (0.45,1.35)
2010	1.92 (1.69,2.19)	2.17 (1.89,2.48)	0.81 (0.47,1.39)
2011	1.62 (1.39,1.89)	1.77 (1.5,2.07)	1.05 (0.63,1.73)
2012	1.61 (1.37,1.9)	1.7 (1.43,2.03)	1.45 (0.93,2.26)
2013	1.09 (0.9,1.31)	1.12 (0.92,1.38)	1.1 (0.67,1.79)
2014	1 (0.82,1.21)	1 (0.81,1.24)	1 (0.62,1.61)


Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B2 on page 99

Figure 5 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, irritant contact dermatitis

a) EPIDERM, all reporters



b) EPIDERM, core reporters



c) EPIDERM, sample reporters




Table 12 Relative risk by year, with 95% comparison intervals, mixed contact dermatitis (2014 estimate = 1)

	Relative risk (95% comparison interval)		
	EPIDERM		
	All reporters	Core reporters	Sample reporters
YEAR			
1996	1.38 (1.12,1.71)	1.51 (1.21,1.89)	0.79 (0.36,1.75)
1997	1.55 (1.29,1.86)	1.66 (1.37,2.01)	1.28 (0.68,2.41)
1998	1.43 (1.19,1.73)	1.55 (1.27,1.89)	1.08 (0.61,1.93)
1999	1.41 (1.17,1.7)	1.51 (1.23,1.85)	1.15 (0.69,1.91)
2000	1.19 (0.97,1.45)	1.33 (1.07,1.64)	0.68 (0.36,1.26)
2001	1.08 (0.88,1.32)	1.25 (1.02,1.54)	0.43 (0.18,1.01)
2002	1.29 (1.08,1.55)	1.45 (1.2,1.74)	0.73 (0.37,1.42)
2003	1.35 (1.13,1.6)	1.55 (1.3,1.86)	0.37 (0.15,0.89)
2004	1.35 (1.14,1.59)	1.49 (1.25,1.77)	0.84 (0.46,1.54)
2005	1.12 (0.93,1.36)	1.26 (1.03,1.54)	0.64 (0.33,1.23)
2006	0.88 (0.71,1.09)	0.96 (0.76,1.2)	0.72 (0.39,1.32)
2007	1.03 (0.84,1.26)	1.2 (0.98,1.47)	0.28 (0.11,0.74)
2008	0.94 (0.76,1.16)	1.06 (0.84,1.32)	0.4 (0.18,0.91)
2009	1.25 (1.03,1.53)	1.45 (1.19,1.78)	0.25 (0.08,0.79)
2010	0.99 (0.79,1.24)	1.02 (0.79,1.31)	1.02 (0.55,1.87)
2011	1.03 (0.8,1.31)	1.06 (0.81,1.38)	0.92 (0.49,1.72)
2012	1.16 (0.92,1.47)	1.22 (0.94,1.57)	0.91 (0.48,1.71)
2013	0.8 (0.61,1.04)	0.8 (0.6,1.07)	0.75 (0.38,1.48)
2014	1 (0.78,1.27)	1 (0.77,1.3)	1 (0.54,1.85)


Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B2 on page 99

Figure 6 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, mixed contact dermatitis

a) EPIDERM, all reporters



b) EPIDERM, core reporters



c) EPIDERM, sample reporters




Table 13 Relative risk by year, with 95% comparison intervals, contact urticaria (2014 estimate = 1)

	Relative risk (95% comparison interval)	
	EPIDERM	
	All reporters	Core reporters
YEAR		
1996	4.57 (3.25,6.43)	5.77 (4.05,8.22)
1997	6.4 (4.98,8.22)	8.05 (6.25,10.36)
1998	4.94 (3.7,6.58)	6.27 (4.68,8.4)
1999	4.99 (3.71,6.72)	5.94 (4.33,8.14)
2000	6.89 (5.39,8.8)	8.87 (6.9,11.4)
2001	4.41 (3.32,5.87)	5.44 (4.05,7.31)
2002	4.94 (3.78,6.45)	6.32 (4.83,8.27)
2003	5.39 (4.17,6.96)	6.9 (5.33,8.93)
2004	4.07 (3.06,5.41)	4.99 (3.72,6.68)
2005	5.76 (4.41,7.52)	7.29 (5.56,9.57)
2006	3.63 (2.61,5.03)	4.02 (2.82,5.72)
2007	3.56 (2.51,5.04)	4.35 (3.04,6.23)
2008	1.95 (1.25,3.05)	2.25 (1.4,3.61)
2009	2.02 (1.3,3.12)	2.32 (1.47,3.67)
2010	3.03 (2.05,4.46)	3.59 (2.4,5.38)
2011	1.84 (1.11,3.06)	2.04 (1.18,3.51)
2012	2.17 (1.35,3.47)	2.59 (1.59,4.21)
2013	1.08 (0.56,2.08)	1.36 (0.71,2.63)
2014	1 (0.53,1.9)	1 (0.49,2.05)


Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model

The number of actual cases on which each analysis is based is provided in Table B2 on page 99

Figure 7 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, contact urticaria

a) EPIDERM, all reporters



b) EPIDERM, core reporters




Table 14 Relative risk by year, with 95% comparison intervals, neoplasia (2014 estimate = 1)

	Relative risk (95% comparison interval)		
	EPIDERM		
	All reporters	Core reporters	Sample reporters
YEAR			
1996	1.96 (1.57,2.45)	4.76 (3.58,6.32)	0.46 (0.18,1.13)
1997	1.92 (1.54,2.39)	4.33 (3.25,5.76)	1.31 (0.73,2.35)
1998	1.54 (1.24,1.92)	3.42 (2.52,4.65)	0.83 (0.45,1.52)
1999	1.61 (1.3,2)	3.87 (2.94,5.1)	0.66 (0.36,1.23)
2000	1.82 (1.49,2.22)	4.15 (3.19,5.4)	0.96 (0.57,1.61)
2001	1.88 (1.57,2.25)	3.98 (3.06,5.17)	1.6 (1.03,2.47)
2002	1.44 (1.18,1.75)	3.09 (2.38,4.02)	1.08 (0.62,1.88)
2003	1.51 (1.25,1.82)	3.31 (2.55,4.29)	1.01 (0.59,1.73)
2004	1.41 (1.16,1.71)	2.91 (2.25,3.78)	1.31 (0.82,2.08)
2005	1.3 (1.06,1.58)	2.95 (2.27,3.83)	0.76 (0.42,1.37)
2006	1.26 (1.04,1.53)	2.4 (1.85,3.12)	1.67 (1.13,2.46)
2007	1.47 (1.2,1.82)	3.02 (2.21,4.13)	1.56 (1.02,2.4)
2008	1.27 (0.98,1.65)	2.79 (2.07,3.76)	0.98 (0.55,1.76)
2009	1.58 (1.24,2.01)	2.93 (2.2,3.89)	1.76 (1.07,2.9)
2010	1.24 (0.9,1.71)	2.2 (1.48,3.28)	1.39 (0.83,2.35)
2011	1.06 (0.63,1.81)	1.27 (0.39,4.09)	1.09 (0.6,1.96)
2012	1.4 (0.88,2.23)	0.99 (0.24,4.12)	1.59 (0.97,2.58)
2013	0.9 (0.56,1.42)	0.8 (0.33,1.93)	1.01 (0.57,1.8)
2014	1 (0.64,1.57)	1 (0.44,2.29)	1 (0.55,1.82)


Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B2 on page 99

Figure 8 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, neoplasia

a) EPIDERM, all reporters



b) EPIDERM, core reporters (Note: scale change)



c) EPIDERM, sample reporters




Table 15 Relative risk by year, with 95% comparison intervals, other (than contact dermatitis) skin (2014 estimate = 1)


	Relative risk (95% comparison interval)		
	EPIDERM		
	All reporters	Core reporters	Sample reporters
YEAR			
1996	3.47 (2.94,4.1)	5.16 (4.38,6.09)	0.89 (0.48,1.67)
1997	3.89 (3.37,4.5)	5.46 (4.66,6.39)	1.82 (1.15,2.88)
1998	3.38 (2.92,3.92)	4.69 (4.01,5.48)	1.6 (1.07,2.39)
1999	3.13 (2.68,3.66)	4.51 (3.83,5.32)	1.31 (0.87,1.97)
2000	3.67 (3.19,4.22)	5.46 (4.71,6.31)	1.16 (0.76,1.77)
2001	3.13 (2.72,3.61)	4.25 (3.66,4.94)	1.79 (1.23,2.6)
2002	2.65 (2.29,3.08)	3.66 (3.14,4.26)	1.32 (0.84,2.07)
2003	2.93 (2.54,3.36)	4.14 (3.59,4.79)	1.13 (0.71,1.79)
2004	2.48 (2.14,2.87)	3.27 (2.79,3.83)	1.65 (1.13,2.39)
2005	2.64 (2.28,3.07)	3.81 (3.26,4.44)	1.03 (0.65,1.63)
2006	2.39 (2.05,2.78)	2.91 (2.45,3.46)	2.19 (1.59,3)
2007	2.52 (2.13,2.97)	3.37 (2.82,4.03)	1.68 (1.14,2.48)
2008	1.75 (1.41,2.18)	2.33 (1.84,2.95)	1.16 (0.7,1.91)
2009	2.09 (1.73,2.52)	2.56 (2.05,3.2)	1.94 (1.26,2.99)
2010	1.92 (1.56,2.37)	2.39 (1.89,3.03)	1.51 (0.94,2.41)
2011	1.28 (0.92,1.79)	1.24 (0.8,1.94)	1.38 (0.84,2.26)
2012	1.65 (1.23,2.22)	1.85 (1.27,2.68)	1.46 (0.91,2.33)
2013	1.11 (0.81,1.53)	1.16 (0.77,1.75)	1.09 (0.65,1.84)
2014	1 (0.72,1.4)	1 (0.65,1.53)	1 (0.58,1.74)

Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model

The number of actual cases on which each analysis is based is provided in Table B2 on page 99

Figure 9 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, other (than contact dermatitis) skin



3.1.6 WORK-RELATED RESPIRATORY DISEASE

The average annual percentage change in risk of work-related respiratory disease, as reported by chest physicians to SWORD is shown in Table 16 whilst the relative rates by year are shown in Tables 17 to 24 and Figures 10 to 17.

The results suggest little change in the annual decrease in incidence of total respiratory disease from -3.0% (95% CIs: -3.7, -2.2) per year (for the period 1999-2013) to -3.2% (95% CIs: -3.9, -2.4) for the current analyses (1999-2014). Similar to the pattern observed for total skin disease, the graphs showing relative rates by year suggest that much of the decrease occurred in the earlier part of the study period (1996-2007 in this instance) with a relatively flat trend until 2013 and a possible decrease between 2013 and 2014. The annual average decrease in the incidence of asthma also remained relatively unchanged from -7.1% (95% CIs: -8.4, -5.9) (for the period 1999-2013) to -7.2% (95% CIs: -8.4, -6.0) for the current analyses (with some suggestion of a decrease between 2013 and 2014). Analyses of shorter-term trends (2007-2014) showed an average change of -4.4% (95% CIs: -8.1, -0.6) per year. An overall decrease in incidence was also observed for all other groups of respiratory disease except pneumoconiosis, for which an annual average increase of 3.2% (95% CIs: 1.1, 5.3) was observed. The graph showing relative rates by year for pneumoconiosis suggests a relatively flat trend in the earlier part of the study period (1999 to 2008) followed by an increasing trend thereafter. For mesothelioma, the data suggest an average annual decrease of -3.0% (95% CIs: -4.3, -1.6) per year, with the annual plots showing a fall in incidence between 2013 and 2014. A smaller annual average decrease was observed for non-malignant pleural disease at -1.2% (95% CIs: -2.3, -0.1) with the annual plots showing a relatively flat trend throughout the study period, with the suggestion of a decrease in incidence between 2013 and 2014. Overall there was little variation by reporter type (core and sample).

Extrapolating the results of the previously described analyses using a zero-inflated binomial model (Section 2.5), the impact of adjusting for 'excess zeros' in the present analyses *might be* expected to change the average annual percentage change in incidence of total work-related respiratory disease in the order of from -3.2% to -2.7%.

Table 16 Average annual percentage change in reported incidence in work-related respiratory disease

		ESTIMATED % CHANGE (95% CONFIDENCE INTERVAL)		
		SWORD		
		All reporters	Core reporters	Sample reporters
	Year (continuous)			
Total respiratory	1999-2014	-3.2 (-3.9, -2.4)	-3.5 (-4.3, -2.6)	-2.4 (-3.7, -1.1)
Asthma	1999-2014	-7.2 (-8.4, -6.0)	-7.0 (-8.3, -5.7)	-8.3 (-11.3, -5.3)
	2007-2014	-4.4 (-8.1, -0.6)	/	/
Mesothelioma	1999-2014	-3.0 (-4.3, -1.6)	-3.1 (-5.0, -1.1)	-2.9 (-4.8, -1.0)
Non-malignant pleural disease	1999-2014	-1.2 (-2.3, -0.1)	-1.9 (-3.2, -0.6)	0.5 (-1.6, 2.6)
• Predominantly plaques	1999-2014	-1.3 (-2.5, 0.0)	-2.2 (-3.7, -0.7)	1.0 (-1.3, 3.4)
• Predominantly diffuse	1999-2014	-0.6 (-2.6, 1.4)	-1.7 (-3.9, 0.5)	4.1 (-0.6, 8.9)
Pneumoconiosis	1999-2014	3.2 (1.1, 5.3)	3.7 (1.1, 6.3)	1.9 (-1.8, 5.7)
	2007-2014	10.4 (4.5, 16.5)	/	/
Other* respiratory disease	1999-2014	-1.5 (-3.0, 0.1)	-1.6 (-3.4, 0.3)	-0.9 (-3.8, 2.2)

*Other than those specified above

Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model

The number of actual cases on which each analysis is based is provided in Table B4 on page 107

Table 17 Relative risk by year, with 95% comparison intervals, total respiratory disease (2014 estimate = 1)

	Relative risk (95% comparison interval)		
	SWORD		
	All reporters	Core reporters	Sample reporters
YEAR			
1999	1.69 (1.55,1.84)	1.78 (1.61,1.97)	1.63 (1.39,1.92)
2000	1.56 (1.43,1.7)	1.7 (1.54,1.88)	1.4 (1.18,1.66)
2001	1.57 (1.44,1.71)	1.77 (1.61,1.95)	1.25 (1.04,1.5)
2002	1.66 (1.52,1.81)	1.92 (1.74,2.12)	1.2 (0.99,1.45)
2003	1.66 (1.52,1.81)	1.99 (1.81,2.18)	1.02 (0.83,1.25)
2004	1.55 (1.42,1.68)	1.83 (1.67,2.01)	1.02 (0.84,1.24)
2005	1.47 (1.35,1.61)	1.64 (1.49,1.81)	1.23 (1.03,1.47)
2006	1.38 (1.26,1.51)	1.5 (1.35,1.66)	1.24 (1.03,1.48)
2007	1.21 (1.09,1.34)	1.25 (1.11,1.42)	1.17 (0.97,1.4)
2008	1.29 (1.17,1.44)	1.36 (1.19,1.55)	1.19 (0.99,1.43)
2009	1.23 (1.1,1.37)	1.32 (1.16,1.51)	1.07 (0.87,1.3)
2010	1.15 (1.03,1.29)	1.27 (1.12,1.45)	0.94 (0.75,1.17)
2011	1.26 (1.12,1.41)	1.35 (1.17,1.55)	1.12 (0.91,1.38)
2012	1.18 (1.05,1.33)	1.3 (1.12,1.49)	1 (0.79,1.25)
2013	1.19 (1.05,1.34)	1.24 (1.06,1.44)	1.12 (0.9,1.38)
2014	1 (0.87,1.14)	1 (0.84,1.19)	1 (0.79,1.26)


Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B4 on page 107

Figure 10 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, total respiratory disease

a) SWORD, all reporters



b) SWORD, core reporters



c) SWORD, sample reporters




Table 18 Relative risk by year, with 95% comparison intervals, asthma (2014 estimate = 1)

	Relative risk (95% comparison interval)		
	SWORD		
	All reporters	Core reporters	Sample reporters
YEAR			
1999	3.18 (2.76,3.67)	3.09 (2.63,3.63)	4.18 (3,5.81)
2000	2.2 (1.87,2.6)	2.11 (1.75,2.55)	2.82 (1.91,4.15)
2001	2.46 (2.1,2.88)	2.58 (2.18,3.05)	1.94 (1.2,3.13)
2002	2.64 (2.24,3.11)	2.78 (2.33,3.32)	2.03 (1.26,3.28)
2003	2.57 (2.18,3.03)	2.77 (2.32,3.29)	1.64 (0.98,2.73)
2004	2.43 (2.04,2.89)	2.65 (2.21,3.18)	1.41 (0.81,2.45)
2005	2.17 (1.81,2.6)	2.26 (1.85,2.75)	1.86 (1.17,2.97)
2006	2.02 (1.7,2.4)	2 (1.65,2.41)	2.45 (1.58,3.79)
2007	1.39 (1.13,1.72)	1.4 (1.11,1.76)	1.44 (0.83,2.49)
2008	1.49 (1.21,1.83)	1.48 (1.18,1.85)	1.75 (1.05,2.9)
2009	1.14 (0.9,1.45)	1.16 (0.9,1.5)	1.09 (0.56,2.11)
2010	1.2 (0.96,1.51)	1.21 (0.95,1.55)	1.28 (0.68,2.39)
2011	1.23 (0.96,1.57)	1.3 (1,1.69)	0.9 (0.43,1.9)
2012	1.28 (1.01,1.61)	1.31 (1.02,1.68)	1.24 (0.62,2.46)
2013	1.25 (0.98,1.59)	1.29 (1,1.67)	1.03 (0.51,2.08)
2014	1 (0.77,1.31)	1 (0.75,1.34)	1 (0.47,2.12)


Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B4 on page 107

Figure 11 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, asthma

a) SWORD, all reporters



b) SWORD, core reporters



c) SWORD, sample reporters




Table 19 Relative risk by year, with 95% comparison intervals, mesothelioma (2014 estimate = 1)

	Relative risk (95% comparison interval)		
	SWORD		
	All reporters	Core reporters	Sample reporters
YEAR			
1999	2.1 (1.79,2.46)	2.54 (2.05,3.15)	2.06 (1.6,2.65)
2000	2.07 (1.77,2.43)	2.78 (2.28,3.39)	1.75 (1.34,2.28)
2001	2.12 (1.81,2.48)	2.76 (2.27,3.37)	1.9 (1.47,2.46)
2002	2.09 (1.78,2.46)	2.82 (2.31,3.45)	1.76 (1.33,2.33)
2003	2.08 (1.77,2.43)	2.91 (2.4,3.54)	1.58 (1.19,2.09)
2004	1.81 (1.54,2.13)	2.6 (2.14,3.15)	1.33 (0.98,1.8)
2005	1.63 (1.37,1.93)	2.2 (1.78,2.73)	1.35 (1.01,1.81)
2006	1.68 (1.39,2.02)	2.42 (1.92,3.05)	1.27 (0.93,1.74)
2007	2.1 (1.73,2.55)	2.65 (2.01,3.5)	1.87 (1.44,2.44)
2008	2.05 (1.67,2.51)	3.43 (2.61,4.49)	1.35 (1,1.83)
2009	1.86 (1.51,2.31)	2.88 (2.18,3.81)	1.31 (0.95,1.82)
2010	1.63 (1.3,2.05)	2.24 (1.64,3.06)	1.36 (0.97,1.89)
2011	1.55 (1.21,1.98)	2.3 (1.65,3.19)	1.17 (0.81,1.69)
2012	1.58 (1.24,2.02)	1.8 (1.25,2.6)	1.58 (1.15,2.18)
2013	1.64 (1.29,2.09)	1.88 (1.29,2.73)	1.61 (1.17,2.2)
2014	1 (0.73,1.36)	1 (0.6,1.67)	1 (0.66,1.52)


Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B4 on page 107

Figure 12 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, mesothelioma

a) SWORD, all reporters



b) SWORD, core reporters



c) SWORD, sample reporters




Table 20 Relative risk by year, with 95% comparison intervals, benign pleural plaques (2014 estimate = 1)

	Relative rates (95% comparison interval)		
	SWORD		
	All reporters	Core reporters	Sample reporters
YEAR			
1999	1.31 (1.14,1.5)	1.42 (1.22,1.65)	1.26 (0.94,1.68)
2000	1.48 (1.3,1.68)	1.68 (1.46,1.93)	1.16 (0.87,1.56)
2001	1.37 (1.21,1.56)	1.64 (1.43,1.89)	0.81 (0.57,1.15)
2002	1.54 (1.36,1.75)	1.84 (1.6,2.11)	0.94 (0.67,1.33)
2003	1.62 (1.44,1.82)	1.93 (1.7,2.18)	0.93 (0.67,1.29)
2004	1.45 (1.29,1.63)	1.68 (1.48,1.91)	0.99 (0.73,1.35)
2005	1.55 (1.39,1.74)	1.69 (1.49,1.92)	1.5 (1.16,1.93)
2006	1.38 (1.22,1.55)	1.39 (1.21,1.6)	1.55 (1.21,1.99)
2007	1.26 (1.09,1.46)	1.34 (1.13,1.59)	1.14 (0.86,1.53)
2008	1.35 (1.16,1.56)	1.33 (1.11,1.6)	1.41 (1.08,1.82)
2009	1.27 (1.08,1.48)	1.32 (1.09,1.59)	1.17 (0.87,1.57)
2010	1.37 (1.17,1.59)	1.5 (1.26,1.79)	1.08 (0.78,1.48)
2011	1.4 (1.19,1.65)	1.47 (1.22,1.78)	1.25 (0.92,1.69)
2012	1.23 (1.04,1.47)	1.31 (1.07,1.6)	1.1 (0.79,1.54)
2013	1.2 (1,1.44)	1.26 (1.01,1.57)	1.07 (0.77,1.49)
2014	1 (0.82,1.23)	1 (0.78,1.28)	1 (0.69,1.44)


Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B4 on page 107

Figure 13 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, benign pleural plaques

a) SWORD, all reporters



b) SWORD, core reporters



c) SWORD, sample reporters




Table 21 Relative risk by year, with 95% comparison intervals, benign pleural plaques – predominantly plaques (2014 estimate = 1)

	Relative risk (95% comparison interval)		
	SWORD		
	All reporters	Core reporters	Sample reporters
YEAR			
1999	1.19 (1.01,1.4)	1.57 (1.32,1.86)	0.57 (0.37,0.88)
2000	1.48 (1.29,1.71)	1.81 (1.54,2.12)	1.07 (0.78,1.47)
2001	1.47 (1.28,1.68)	1.88 (1.62,2.18)	0.85 (0.59,1.21)
2002	1.59 (1.38,1.84)	2.04 (1.76,2.38)	0.88 (0.61,1.27)
2003	1.7 (1.5,1.94)	2.17 (1.89,2.49)	0.95 (0.67,1.34)
2004	1.48 (1.3,1.68)	1.92 (1.67,2.21)	0.68 (0.47,1)
2005	1.65 (1.46,1.86)	1.95 (1.71,2.24)	1.37 (1.04,1.8)
2006	1.28 (1.11,1.48)	1.43 (1.21,1.68)	1.19 (0.89,1.59)
2007	1.22 (1.04,1.44)	1.45 (1.21,1.75)	0.88 (0.63,1.23)
2008	1.28 (1.08,1.52)	1.48 (1.21,1.81)	0.95 (0.68,1.33)
2009	0.97 (0.79,1.18)	1.27 (1.02,1.59)	0.43 (0.26,0.71)
2010	1.35 (1.13,1.61)	1.57 (1.28,1.93)	1 (0.7,1.42)
2011	1.34 (1.11,1.62)	1.58 (1.26,1.97)	0.92 (0.64,1.33)
2012	1.3 (1.07,1.58)	1.47 (1.16,1.86)	1.06 (0.74,1.52)
2013	1.26 (1.02,1.54)	1.4 (1.09,1.8)	1.02 (0.72,1.46)
2014	1 (0.79,1.27)	1 (0.74,1.36)	1 (0.68,1.47)


Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B4 on page 107

Figure 14 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, benign pleural plaques – predominantly plaques

a) SWORD, all reporters



b) SWORD, core reporters



c) SWORD, sample reporters




Table 22 Relative risk by year, with 95% comparison intervals, benign pleural plaques – predominantly diffuse (2014 estimate = 1)

	Relative risk (95% comparison interval)		
	SWORD		
	All reporters	Core reporters	Sample reporters
YEAR			
1999	1.07 (0.83,1.38)	1.17 (0.89,1.52)	0.82 (0.34,1.98)
2000	1.35 (1.08,1.7)	1.54 (1.22,1.94)	0.75 (0.31,1.82)
2001	1.18 (0.93,1.5)	1.34 (1.05,1.71)	0.61 (0.23,1.64)
2002	1.49 (1.19,1.87)	1.64 (1.3,2.07)	1.24 (0.59,2.61)
2003	1.43 (1.15,1.79)	1.61 (1.28,2.02)	0.93 (0.42,2.08)
2004	1.03 (0.81,1.33)	1.19 (0.92,1.53)	0.43 (0.14,1.33)
2005	1.24 (0.98,1.56)	1.26 (0.98,1.61)	1.92 (1.07,3.43)
2006	1.46 (1.16,1.84)	1.37 (1.05,1.77)	2.59 (1.59,4.21)
2007	1.27 (0.95,1.7)	1.13 (0.8,1.59)	2.05 (1.18,3.56)
2008	1.32 (0.99,1.77)	1.16 (0.82,1.63)	2.14 (1.26,3.63)
2009	1.39 (1.04,1.85)	1.19 (0.85,1.67)	2.44 (1.44,4.15)
2010	1.31 (0.98,1.76)	1.47 (1.08,1.99)	0.49 (0.16,1.54)
2011	1.35 (0.99,1.84)	1.27 (0.89,1.8)	1.82 (0.94,3.5)
2012	0.93 (0.65,1.33)	0.98 (0.67,1.43)	0.67 (0.22,2.08)
2013	1.13 (0.8,1.59)	1.07 (0.72,1.58)	1.44 (0.71,2.95)
2014	1 (0.69,1.44)	1 (0.67,1.5)	1 (0.41,2.41)


Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B4 on page 107

Figure 15 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, benign pleural plaques – predominantly diffuse


a) SWORD, all reporters



b) SWORD, core reporters



c) SWORD, sample reporters (note scale change)



**Table 23 Relative risk by year, with 95% comparison intervals,
pneumoconiosis (2014 estimate = 1)**

	Relative risk (95% comparison interval)		
	SWORD		
	All reporters	Core reporters	Sample reporters
YEAR			
1999	0.62 (0.49,0.79)	0.57 (0.43,0.75)	0.88 (0.56,1.37)
2000	0.59 (0.47,0.75)	0.62 (0.47,0.8)	0.55 (0.31,0.96)
2001	0.54 (0.42,0.69)	0.51 (0.38,0.69)	0.67 (0.4,1.12)
2002	0.51 (0.39,0.67)	0.58 (0.43,0.77)	0.28 (0.13,0.62)
2003	0.55 (0.42,0.7)	0.61 (0.47,0.81)	0.32 (0.15,0.67)
2004	0.45 (0.34,0.59)	0.53 (0.4,0.71)	0.16 (0.06,0.42)
2005	0.54 (0.42,0.69)	0.58 (0.44,0.76)	0.48 (0.27,0.86)
2006	0.61 (0.48,0.77)	0.67 (0.52,0.88)	0.46 (0.25,0.83)
2007	0.41 (0.29,0.58)	0.39 (0.26,0.59)	0.48 (0.27,0.85)
2008	0.54 (0.39,0.73)	0.56 (0.39,0.81)	0.5 (0.28,0.91)
2009	0.84 (0.65,1.09)	0.98 (0.73,1.31)	0.49 (0.26,0.91)
2010	0.54 (0.4,0.74)	0.64 (0.46,0.9)	0.3 (0.14,0.67)
2011	0.8 (0.6,1.05)	0.86 (0.62,1.18)	0.69 (0.4,1.18)
2012	0.66 (0.48,0.89)	0.72 (0.51,1.02)	0.53 (0.27,1.02)
2013	0.93 (0.71,1.22)	0.99 (0.72,1.36)	0.83 (0.51,1.36)
2014	1 (0.76,1.31)	1 (0.72,1.39)	1 (0.62,1.62)


Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B4 on page 107

Figure 16 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, pneumoconiosis

a) SWORD, all reporters



b) SWORD, core reporters



c) SWORD, sample reporters




Table 24 Relative risk by year, with 95% comparison intervals, other (than those investigated separately) respiratory disease (2014 estimate = 1)

	Relative risk (95% comparison interval)		
	SWORD		
	All reporters	Core reporters	Sample reporters
YEAR			
1999	1.19 (0.94,1.49)	1.24 (0.96,1.61)	0.99 (0.61,1.59)
2000	1.15 (0.93,1.44)	1.08 (0.83,1.4)	1.37 (0.91,2.07)
2001	1.28 (1.04,1.59)	1.24 (0.96,1.59)	1.39 (0.92,2.1)
2002	1.33 (1.07,1.66)	1.41 (1.1,1.82)	1.02 (0.64,1.64)
2003	1.27 (1.03,1.57)	1.34 (1.06,1.69)	1.09 (0.69,1.74)
2004	1.46 (1.2,1.78)	1.58 (1.27,1.97)	1.13 (0.73,1.76)
2005	0.87 (0.68,1.1)	0.88 (0.66,1.15)	0.88 (0.53,1.47)
2006	1.16 (0.93,1.43)	1.25 (0.99,1.58)	0.85 (0.51,1.41)
2007	0.84 (0.65,1.09)	0.85 (0.63,1.15)	0.79 (0.47,1.32)
2008	1.16 (0.91,1.47)	1.22 (0.93,1.6)	0.99 (0.61,1.61)
2009	1.31 (1.05,1.64)	1.29 (0.99,1.68)	1.34 (0.88,2.04)
2010	0.88 (0.67,1.16)	1 (0.75,1.35)	0.52 (0.26,1.04)
2011	1.08 (0.83,1.4)	0.89 (0.63,1.25)	1.68 (1.1,2.56)
2012	1.16 (0.89,1.51)	1.34 (1.01,1.79)	0.65 (0.33,1.26)
2013	1.07 (0.82,1.41)	0.93 (0.66,1.33)	1.36 (0.87,2.11)
2014	1 (0.75,1.33)	1 (0.71,1.4)	1 (0.56,1.77)


Models adjusted for reporter type (where appropriate), season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B4 on page 107

Figure 17 Relative risk by year (2014 estimate = 1), with 95% comparison intervals, other (than those investigated separately) respiratory disease


a) SWORD, all reporters



b) SWORD, core reporters



c) SWORD, sample reporters



3.1.7 WORK-RELATED MUSCULOSKELETAL DISORDERS

The average annual percentage change in reported incidence of work-related musculoskeletal disorders (MSDs), as reported by GPs (THOR-GP) is shown in Table 25 whilst the relative rates by year are shown in Tables 26 to 29 and Figures 18 to 21. Data from GPs reporting as core reporters suggested a downward trend in the incidence of total work-related MSDs in the order of -15.8% (95% CIs: -18.9, -12.1) per year for the period 2006-2009. The graph showing relative rates by year suggested an overall downward trend, with the largest decrease being between 2006 and 2007. A similar annual pattern was seen for the subset of upper limb disorders, with an average annual decrease of -14.7% (95% CIs: -19.6, -9.5). A steeper decrease was observed for spine/back disorders at -19.7% (95% CIs: -24.9, -14.1) with the annual plots suggesting a continual decrease throughout the study period, whilst for lower limb disorders an initial decrease between 2006 and 2007 was followed by a relatively flat trend: average annual decrease for this group of -10% (95% CIs: -19.5, +0.6) for lower limb disorders.

For sample reporters, the annual average decrease in incidence for total musculoskeletal disorders (2011-2014) was -12.1% (95% CIs: -21.3, -1.7). This compared to -4.5% (95% CIs: -21.1, +15.5) reported previously (based on data for 2011 to 2013). The graph showing relative rates by year suggest a relatively flat trend between 2011 and 2013, followed by a decrease between 2013 and 2014 (although confidence intervals were wide and overlapping). Some variation was observed between the different MSD sub-groups but confidence intervals were very wide. For upper limb disorders an annual average decrease of -1.8% (95% CIs: -16.5, 15.3) was observed whilst for spine/back the equivalent figure was -17.3% (95% CIs: -30.3, -1.9) and for lower limb it was -8.8% (95% CIs: -30.6, 19.9).

Table 25 Average annual percentage change in reported incidence in total work-related musculoskeletal disorders

		ESTIMATED % CHANGE (95% CONFIDENCE INTERVAL)	
		THOR-GP	
		Core reporters	Sample reporters
	Year (continuous)		
Total musculoskeletal	2006-2009	-15.8 (-19.4, -12.1)	/
	2011-2014	/	-12.1 (-21.3, -1.7)
Upper limb	2006-2009	-14.7 (-19.6, -9.5)	
	2011-2014	/	-1.8 (-16.5, 15.3)
Spine/back	2006-2009	-19.7 (-24.9, -14.1)	/
	2011-2014	/	-17.3 (-30.3, -1.9)
Lower limb	2006-2009	-10 (-19.5, 0.6)	/
	2011-2014	/	-8.8 (-30.6, 19.9)

Models adjusted for, season and harvesting

Population offset included in the model

The number of actual cases on which each analysis is based is provided in Table B6 on page 117

Table 26 Relative risk by year, with 95% comparison intervals, total work-related musculoskeletal disorders (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1)

Relative risk (95% comparison interval)		
THOR-GP		
	Core reporters	Sample reporters
YEAR		
2006	1.75 (1.61,1.92)	/
2007	1.27 (1.18,1.38)	/
2008	1.19 (1.1,1.3)	/
2009	1 (0.9,1.11)	/
2010	/	/
2011	/	1.44 (1.17,1.76)
2012	/	1.4 (1.16,1.7)
2013	/	1.33 (1.09,1.63)
2014	/	1 (0.79,1.26)


Models adjusted for, season and harvesting

Population offset included in the model

The number of actual cases on which each analysis is based is provided in Table B6 on page 117

Figure 18 Relative risk by year (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1), with 95% comparison intervals, total musculoskeletal disorders

a) THOR-GP, core reporters



b) THOR-GP, sample reporters




Table 27 **Relative risk by year, with 95% comparison intervals, upper limb disorders (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1)**

	Relative risk (95% comparison interval)	
	THOR-GP	
	Core reporters	Sample reporters
YEAR		
2006	1.63 (1.44,1.84)	/
2007	1.21 (1.08,1.35)	/
2008	1.07 (0.95,1.2)	/
2009	1 (0.87,1.15)	/
2010	/	/
2011	/	1.17 (0.86,1.6)
2012	/	1.1 (0.81,1.5)
2013	/	1.34 (1.01,1.77)
2014	/	1 (0.72,1.4)


Models adjusted for season and harvesting

Population offset included in the model

The number of actual cases on which each analysis is based is provided in Table B6 on page 117

Figure 19 Relative risk by year (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1), with 95% comparison intervals, upper limb disorders

a) THOR-GP, core reporters



b) THOR-GP, sample reporters




Table 28 Relative risk by year, with 95% comparison intervals, spine/back disorders (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1)

	Relative risk (95% comparison interval)	
	THOR-GP	
	Core reporters	Sample reporters
YEAR		
2006	2.11 (1.84,2.42)	/
2007	1.56 (1.38,1.77)	/
2008	1.44 (1.26,1.64)	/
2009	1 (0.85,1.18)	/
2010	/	/
2011	/	1.68 (1.25,2.25)
2012	/	1.76 (1.32,2.35)
2013	/	1.54 (1.13,2.1)
2014	/	1 (0.68,1.47)


Models adjusted for season and harvesting

Population offset included in the model

The number of actual cases on which each analysis is based is provided in Table B6 on page 117

Figure 20 Relative risk by year (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1), with 95% comparison intervals, spine/back disorders

a) THOR-GP, core reporters



b) THOR-GP, sample reporters




Table 29 **Relative risk by year, with 95% comparison intervals, lower limb disorders (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1)**

	Relative risk (95% comparison interval)	
	THOR-GP	
	Core reporters	Sample reporters
YEAR		
2006	1.51 (1.19,1.92)	/
2007	0.95 (0.75,1.2)	/
2008	1.14 (0.92,1.41)	/
2009	1 (0.78,1.28)	/
2010	/	/
2011	/	1.09 (0.68,1.74)
2012	/	0.96 (0.57,1.6)
2013	/	0.68 (0.38,1.2)
2014	/	1 (0.59,1.68)


Models adjusted for season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B6 on page 117

Figure 21 Relative risk by year (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1), with 95% comparison intervals, lower limb disorders

a) THOR-GP, core reporters



b) THOR-GP sample reporters



3.1.8 WORK-RELATED MENTAL ILL-HEALTH

The average annual percentage change in reported incidence of work-related mental ill-health, as reported by GPs (THOR-GP) is shown in Table 30 whilst the relative rates by year are shown in Tables 31 to 33 and Figures 22 to 24.

Based on data from core reporters, an average annual decrease in the incidence of total mental ill-health of -12.4% (95% CIs: -17.1, -7.4) was observed (2006-2009). This compared to an average annual decrease of -11.4% (95% CIs: -18.6, -3.7) for anxiety and depression and -13.3% (95% CIs: -19, -7.1) for other work stress. Overall, the graphs showing relative rates by year suggest a general decrease in incidence over the study period.

For sample reporters, the annual average decrease in incidence for total mental ill-health (2011-2014) was -15.0% (95% CIs: -25.1, -3.5). This compared to -6.7% (95% CIs: -24.2, 14.9) reported previously (based on data for 2011 to 2013). The annual plot suggests a flat trend between 2011 and 2012, which decreases thereafter (although confidence intervals were wide and overlapping). The equivalent changes in incidence for anxiety and depression and other work stress were -20.6% (95% CIs: -36.6, -0.5) and -9.7% (95% CIs: -22.2, 4.7), respectively.

Table 30 Average annual percentage change in reported incidence in total work-related mental ill-health

		ESTIMATED % CHANGE (95% CONFIDENCE INTERVAL)	
		THOR-GP	
		Core reporters	Sample reporters
	Year (continuous)		
Total mental ill-health	2006-2009	-12.4 (-17.1, -7.4)	/
	2011-2014	/	-15.0 (-25.1, -3.5)
Anxiety and depression	2006-2009	-11.4 (-18.6, -3.7)	/
	2011-2014	/	-20.6 (-36.6, -0.5)
Other work stress	2006-2009	-13.3 (-19, -7.1)	/
	2011-2014	/	-9.7 (-22.2, 4.7)

Models adjusted for season and harvesting

Population offset included in the model

The number of actual cases on which each analysis is based is provided in Table B6 on page 117

Table 31 **Relative risk by year, with 95% comparison intervals, total mental ill-health (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate =1)**

	Relative risk (95% comparison interval)	
	THOR-GP	
	Core reporters	Sample reporters
YEAR		
2006	1.54 (1.37,1.74)	/
2007	1.4 (1.27,1.55)	/
2008	1.27 (1.15,1.41)	/
2009	1 (0.88,1.14)	/
2010	/	/
2011	/	1.56 (1.25,1.95)
2012	/	1.51 (1.22,1.88)
2013	/	1.32 (1.06,1.65)
2014	/	1 (0.77,1.31)


Models adjusted for season and harvesting

Population offset included in the model

The number of actual cases on which each analysis is based is provided in Table B6 on page 117

Figure 22 Relative risk by year (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1), with 95% comparison intervals, total mental ill-health

a) THOR-GP, core reporters



b) THOR-GP, sample reporters




Table 32 **Relative risk by year, with 95% comparison intervals, anxiety and depression (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1)**

	Relative risk (95% comparison interval)	
	THOR-GP	
	Core reporters	Sample reporters
YEAR		
2006	1.4 (1.17,1.68)	/
2007	1.35 (1.17,1.57)	/
2008	1.12 (0.95,1.32)	/
2009	1 (0.82,1.21)	/
2010	/	/
2011	/	1.76 (1.18,2.61)
2012	/	1.71 (1.17,2.48)
2013	/	1.23 (0.8,1.89)
2014	/	1 (0.61,1.63)


Models adjusted for season and harvesting

Population offset included in the model

The number of actual cases on which each analysis is based is provided in Table B6 on page 117

Figure 23 Relative risk by year (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1), with 95% comparison intervals, anxiety and depression

a) THOR-GP, core reporters



b) THOR-GP, sample reporters (note scale change)




Table 33 Relative rates by year, with 95% comparison intervals, other work stress (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1)

	Relative rates (95% comparison interval)	
	THOR-GP	
	Core reporters	Sample reporters
YEAR		
2006	1.71 (1.47,1.98)	/
2007	1.48 (1.3,1.68)	/
2008	1.47 (1.3,1.67)	/
2009	1 (0.85,1.18)	/
2010	/	/
2011	/	1.43 (1.1,1.85)
2012	/	1.2 (0.91,1.59)
2013	/	1.27 (0.97,1.65)
2014	/	1 (0.74,1.36)


Models adjusted for season and harvesting

Population offset included in the model


The number of actual cases on which each analysis is based is provided in Table B6 on page 117

Figure 24 Relative risk by year (analyses based on core reporters 2009 estimate = 1, analyses based on sample reporters 2014 estimate = 1), with 95% comparison intervals, other work related stress

a) THOR-GP, core reporters



b) THOR-GP, sample reporters



4 DISCUSSION

This report describes temporal trends in incidence of WRI in the UK as reported to the three constituent schemes of THOR which are currently funded by HSE. These are EPIDERM (dermatologists), SWORD (chest physicians) and THOR-GP (GPs). It updates on previously submitted reports³⁻¹¹ by the incorporation of a further year (2014) of data. The method employed has been described in full in both the current and preceding reports. Essentially, a longitudinal, negative binomial (i.e. over-dispersed) Poisson model with random effects was fit to the data. This enabled change over time in the number of reporters and in other reporter characteristics which could independently impact on case density to be taken into account.

As agreed with HSE, the trend estimates presented here have not yet been formally adjusted for the impact of 'reporter fatigue'. The extensive body of work undertaken to investigate whether THOR reporters are exhibiting reporter fatigue is described in Section 2.5 and in previous reports^{3, 6, 12-15}. Most recently these analyses have focused on whether fatigue may be manifesting as an excess of zero reports in the data, and whether the proportion of 'excess zeros' has increased the longer a reporter has participated in the scheme¹⁶. The results of these investigations have suggested that for both EPIDERM and SWORD, there is some evidence of fatigue manifesting in this way but that the magnitude is different for the two schemes and tended to be greater for sample compared to core reporters. There was also some evidence of fatigue amongst GPs reporting to THOR-GP. Here though, the effect appeared to be greater for core reporters, with little evidence of a significant increase in excess zeros over time for sample reporters who comprise the currently reporting participants.

As previously agreed with HSE, the trend estimates presented in the annual reports will not be formally adjusted for fatigue until after the methodology has been through the peer review process (which is anticipated to be completed before the 2016 trends report deadline). However, as in the report submitted in 2014, we have endeavoured to provide an estimate as to the *possible* impact of adjusting for fatigue on the current trend results. It is important to stress though that these are estimates, provided for guidance purposes only. Differences between the current datasets and those used in the zero-inflated binomial models (which only included data up to 2012) and differences in the modelling procedure means that applying the zero-inflated binomial model to the current data would likely yield different results to the estimates provided here.

An abridged commentary by category of illness is provided in the following sections.

SKIN (EPIDERM and THOR-GP): The primary THOR data source on work-related skin disease is EPIDERM with approximately 18000 case reports during the study period (1996-2014). Trends based on this data source have been reported annually

to HSE since 2004 and the estimated average annual change in incidence has remained largely unchanged by the addition of each successive year of data, at between 3-4%. However, the annual plots do suggest some variation from year to year. Overall, there appears to have been a general decrease in incidence in the earlier part of the study period (1996 to 2007), followed by a levelling out between 2007 and 2012 and then a further drop between 2012 and 2013. There was little suggestion of the incidence continuing to fall between 2013 and 2014. Although reporter fatigue (exhibiting as an increase in zeros over membership time) is likely present in both EPIDERM core and sample reporters, it appears to be more extensive in the latter, perhaps because they are less committed to the scheme or have less sophisticated systems than the core reporters who tend to have a strong interest in the area and who tend to work in larger referral centres. However, because sample reporters contribute less data overall compared to core reporters (12%), the *likely* impact on the overall estimate may be relatively small (remaining at a 3-4% decrease in incidence per year).

Work-related skin disease is also currently reported by GPs to THOR-GP and by occupational physicians to OPRA (not reported upon here). Trends based on GP data are either based on core reports only (2006-2009) or sample reports only (2011-2014). The former have been presented previously (and are again included here for completeness) and suggest a bigger annual decrease (compared EPIDERM trends based on core data for the same period) of 6%. Trends based on GP sample only data (2011-2014) suggest an average, annual decrease of approximately 19%. However, it should be noted that the confidence intervals for this estimate were wide and included zero. Similarly, it is not possible to discern any pattern from the annual plots (confidence intervals are wide and overlapping).

The impact of reporter fatigue on GP reported skin disease trends cannot be directly estimated. There is evidence that core reporters (who reported during 2006 and 2010) experienced fatigue, and if so, the trend estimates for this group would be attenuated. However, the fatigue analyses were based on total cases and cannot be directly extrapolated to specific disease (e.g. skin) groups. There was no evidence that the proportion of excess zeros increased over time for THOR-GP sample reporters.

The majority (82%) of the case reports to EPIDERM were CD and therefore unsurprisingly the observed trends for this group were very similar to those observed for total skin disease i.e. an average annual decrease in CD incidence of -3.7% (compared to -3.8 for total skin) and an earlier decrease in incidence followed (circa 2006) by a relatively flat trend until a further drop in 2013. As discussed previously, in addition to investigating CD trends overall, the MLM methodology (or an adaptation of) has also been applied to investigate changes in incidence of CD related to specific agents or economic sectors²²⁻²⁶. In doing so, we have shown a reduction in incidence of CD in cement workers attributed to chromate, and in

healthcare workers attributed to latex (in response to specific Government interventions aimed at reducing exposure to these agents)^{22, 23} but also an increase in incidence of CD in nail technicians attributed to acrylates, and in healthcare workers attributed to methylchloroisothiazolinone/methylisothiazolinone (MCI/MI) and to handwashing²⁴⁻²⁶. Thus, whilst the incidence of dermatologist reported CD may be falling overall, the extent to which it is falling may vary between workers and for certain groups with specific exposures, it may even be increasing.

Dermatologist reported trends for the different categories of CD (allergic, irritant and mixed allergic/irritant) were also investigated, the results of which suggest a larger overall decrease in incidence observed for allergic compared to irritant (or mixed) CD. This disparity probably reflects the aforementioned Government interventions (UK/EU) aimed at reducing allergic CD attributed to specific agents (latex, chromate). However, of interest, the drop in incidence between 2012 and 2013 (observed for total CD) appears to be largely driven by a drop in irritant rather than allergic CD between these two years. It remains to be seen whether this apparent trend is maintained so it may be premature to speculate on the explanation for this observation.

A statistically significant annual average decrease in incidence was also observed for dermatologist reported (to EPIDERM) contact urticaria and neoplasia. For neoplasia this was of a similar order to that observed for CD (approximately 3% per year) whilst for urticaria it was larger at approximately 7% per year. It has been postulated previously that the overall decline in neoplasia case reports to EPIDERM likely reflects a decline in the number of cases arising from sun exposure (of armed forces). However, consultation with key dermatologists at the 2015 EPIDERM Advisory Meeting suggested it may also reflect the fact that dermatologists currently reporting to EPIDERM (particularly core reporters) largely specialise in CD and therefore these neoplasia cases are not being captured. To increase the representativeness of EPIDERM for this diagnostic group may therefore require targeted recruitment of those dermatologists specialising in neoplasia. This is being explored.

For both neoplasia and urticaria, markedly different trends were observed for core and sample reporters, with core data suggesting a decrease in incidence and sample data suggesting a much smaller decrease (urticaria) or an increase (neoplasia) in incidence. Furthermore, if EPIDERM sample reporters are experiencing greater fatigue than core reporters (shown for total skin disease) then the disparity between the core and sample trend estimates may become even larger. However, for neoplasia, the confidence intervals for the annual plots are wide and overlapping for both core and sample reporters, again suggesting that EPIDERM in general may not be particularly capturing these cases.

It is also useful to compare trends derived from THOR data with data from other, external sources. For example, as part of the work undertaken by the Modernet group (an EU wide network for development of new techniques for discovering trends in WRI and tracing new and emerging risks), trends in physician reported CD have been compared across 10 European countries²⁷. The results showed a similarity in CD trends across the different countries, with data for most countries suggesting a decline in incidence. THOR data can also be compared with data from the Self-reported Work-related Illness (SWI) survey, conducted annually as part of the Labour Force Survey (LFS)²⁸. The latest 3-year estimates (numbers are typically too small to provide reliable annual estimates) from the SWI are for the period averaged 2010/11, 2011/12, 2013/14 and suggest a decline in incidence compared to the previous period (2009/10-2011/12) from 22 per 100,000 employed to 17 per 100,000 employed²⁹.

RESPIRATORY (SWORD): Chest physicians reporting to SWORD are the primary THOR source of case reports of work-related respiratory disease with both OPs and GPs reporting relatively few respiratory diagnoses (<5% of total cases reported by these two groups). Trends based on data from chest physicians have also been reported since the initial report submitted to HSE in 2006³ but unlike dermatologists, the addition of each successive year of data appears to have had more of an impact on the trend estimate (from an initial 1% annual decrease to the 3% currently observed). It was previously suggested that this probably reflects the fact that compared to EPIDERM (where reports are predominantly of CD and neoplasia, and have been throughout the study period), case reports to SWORD encompass a wider diagnostic range with the proportion of the total cases attributed to each diagnosis exhibiting some variation throughout the study period.

Overall, the likely impact of fatigue on the trend estimate (for total respiratory disease) is relatively small (a possible reduction in the annual decrease from approximately -3.2% to -2.7%). Whilst there is evidence that SWORD sample reporters are exhibiting fatigue (manifesting as an increase in zero cases reports over membership time) there is little evidence of the same phenomenon amongst SWORD core reporters (probably reflecting the strong commitment of stalwart 'core' SWORD reporters).

With the addition of the 2014 data, the estimated annual average percentage change in reported incidence of asthma has remained at approximately 7% per year. The annual plots suggest a similar pattern as that observed for total skin disease with a decrease in incidence in the earlier part of the study followed by a relatively flat trend (in this case since approximately 2007). There is a slight suggestion of a further decrease between 2013 and 2014 but confidence intervals for the individual years are overlapping. As for skin disease, it is important to view these 'overall' changes in incidence in conjunction with the results from other studies investigating changes in

incidence of WRI related to specific agents, Government interventions etc. For example, Stocks *et al*³⁰⁻³¹ observed a significant reduction in reports of asthma attributed to agents with a work exposure limit (WEL) relative to those without a WEL²⁹. Of interest, however, for some agents, for example flour, a significant increase in the incidence of asthma (relative to other agents) was observed³⁰. This is disappointing to note especially in view of longstanding attempts at dissemination of knowledge of asthma risks associated with flour and other substances involved in baking.

As described for skin disease, it is again of use to compare SWORD trends with other data both within the UK and the EU. The 3-year average SWI derived incidence rate for 'breathing or lung problems' suggest a decline in incidence from 42 per 100,000 employed in 2009/10-2011/12 to 34 per 100,000 employed in 2010/11, 2011/12, 2013/14²⁹ whilst results from Modernet again suggest similarities across the participating countries, with an overall decline in the incidence of asthma²⁷.

The other main groups of respiratory disease reported to SWORD are the (primarily) asbestos related diseases, namely, mesothelioma, benign pleural plaques and pneumoconiosis. For mesothelioma, the results suggest an annual average decrease of approximately 3% per year (compared to 2% per year reported previously). The annual plot show a relatively flat trend since approximately 2010 but there is suggestion of a significant fall in incidence between 2013 and 2014 (which is particularly pronounced in the core reports). This decreasing or flat trend is contrary to what is expected as other evidence (including epidemiological studies from Peto *et al* and the mesothelioma death registers) suggests the incidence is rising with a possible peak, expected in 2016^{32, 33}. Consultation with key SWORD reporters suggested that the decline in case reports of mesothelioma to SWORD probably reflects a shift in referral patterns since a proportion of these cases are now being referred (via the 2 week rapid access pathway) to oncologists (or to chest physicians specialising in lung cancer who may not report to THOR) rather than exclusively to chest physicians, as used to be the case. This issue continues to be discussed at the annual meetings of the SWORD Advisory Committee, with one suggestion being that THOR could approach lung cancer specialists and/or possibly the non-specialist physicians who organise and run the rapid access systems and ask them to report to SWORD.

A relatively flat trend was also observed for benign pleural plaques although there was again a suggestion of a decrease in incidence between 2013 and 2014. As discussed previously, this probably reflects the fact that individuals presenting with this abnormality alone (in England and Wales) are no longer financially compensated³⁴ and therefore, referrals to chest physicians are less common. This was further corroborated by data for Scotland (where compensation is still available) for which analysis of the 'crude' data suggested an increase in case reports over time.

Data from SWORD continue to suggest a possible increase in pneumoconiosis incidence since approximately 2007. Whilst the majority (77%) of case reports of pneumoconiosis during the study period were attributed to asbestos, a significant proportion was attributed to other agents, for example silica (11%) and coal (8%). As such, work is ongoing to investigate whether differing trends would be apparent if trends were investigated separately for specific agents.

MUSCULOSKELETAL AND MENTAL ILL-HEALTH (THOR-GP): THOR-GP currently provides the only (HSE funded) THOR source of data on work-related MSDs and mental ill-health. Trends for these two disease groups have also been reported previously for occupational physicians reporting to OPRA, rheumatologists reporting to MOSS and psychiatrists reporting to SOSMI^{7, 8}. Because of the well documented extensive change from predominantly core to 100% sample reporting (and the resulting possible impact on incidence)¹⁹, trends based on THOR-GP data are presented separately for core (2006-2009) and sample (2011 onwards) only. The results of analyses based on the former have been reported previously but have again been included here for completeness. Overall, the results suggest an annual average decrease in the incidence of GP reported MSDs of approximately 16% per year. Restricting the analyses to specific MSD sub-groups, a steeper decrease was observed for spine/back disorders (approximately 20% per year) compared to upper limb disorders (15% per year). An overall decrease in mental ill-health incidence was also observed for this group, with similar estimates for total cases and the two subgroups of anxiety and depression and work stress (11-13%). As discussed, there was some evidence that GP core reporters were experiencing fatigue (manifesting as an increase in excess zeros over time), which if true (and if applicable across the different disease groups) would mean these trends would be attenuated slightly.

However, since core reporting has now ceased it is trends based on sample data that are more informative for current and future trends. These data also suggest an overall decrease in incidence of work-related MSDs, which is of a similar magnitude (12%) to that observed for core reporters over the earlier period (16%). This estimated annual decrease (12%) is considerably larger to that reported previously (5%) and appears to be largely driven by an apparent decrease in incidence between 2013 and 2014 (although confidence intervals are overlapping). This is also the first report to present sample only trends for the MSD sub-groups of upper limb, spine/back and lower limb disorders. For spine/back disorders the annual average decrease (17%) was comparable with that observed for core reporters for the earlier period (20%). For the other two diagnostic groups (upper limb and lower limb) the confidence intervals were very wide making it difficult to draw any firm conclusions at this stage.

Sample data from GPs (2011-2014) also suggested an overall downward trend in the incidence of work-related mental ill-health (which at 15% is similar to the 12%

observed for the core reporters during the earlier period) which is bigger than that reported previously (although confidence intervals for the estimate reported in 2014 are wide). A large, significant annual decrease was also observed for anxiety and depression (which was of a similar magnitude to that reported previously). As discussed, investigations of fatigue have not suggested any evidence of this phenomenon for GP sample reporters. The other main dataset collecting data on work-related musculoskeletal and mental ill-health in the UK (for comparison with THOR data) is the SWI. Unlike THOR-GP data, data from the SWI suggest little change in incidence of work-related musculoskeletal disorders from 540 per 100,000 employed in 2009/10-2011/12 to 530 per 100,000 employed in 2010/11, 2011/12, 2013/14²⁹. In comparison, SWI data for 'stress, depression or anxiety' suggest no change at 740 per 100,000 employed for both time periods²⁹.

5. CONCLUSIONS

Case reports to THOR have been utilised to investigate trends in incidence of medically reported WRI since the initial report to HSE in 2005. In addition to the main reports (submitted annually) which have not only updated the trends but have also described ongoing methodological developments, additional reports have been submitted describing further investigations into important issues such as reporter fatigue. Whilst overall the observed trends have remained relatively unchanged with the addition of each successive year of data and are in accordance with those expected as a result of Government initiatives (for example, the general decline in incidence of asthma, contact dermatitis and contact urticaria) others have shown more variation (for example, the asbestos related diseases). Furthermore, trends related to specific sectors or agents often appear discordant with the 'overall' trends (for example asthma and flour or CD and handwashing), thus showing the value of THOR in identifying real and significant adverse or desired trends in relation to specific exposures. Ongoing work, including benchmarking with other data sources and investigating trends in relation to specific sectors or agents will help clarify these issues further.

ACKNOWLEDGEMENTS

THOR is partially funded by the Health and Safety Executive and has also received funding from other sources. We are grateful to the physicians who report to THOR for their continuing support. The opinions expressed in this report are of the researchers and not necessarily those of the HSE.

REFERENCES

1. McNamee R, Carder M, Chen Y and Agius R. Assessment of changes in the incidence of work-related disease over time using ODIN and THOR surveillance data. Report to HSE submitted 2005
2. The THOR website. Available at: <http://www.coeh.man.ac.uk/thor> (last accessed August 2015)
3. McNamee R, Carder M, Chen Y, Agius R. Time trends in the incidence of work-related disease in the UK, 1996-2004: estimation from ODIN/THOR surveillance data. Report to HSE submitted September 2006.
4. McNamee R, Carder M, Chen Y, Agius R. Time trends in the incidence of work-related disease in the UK, 1996-2006: estimation from ODIN/THOR surveillance data. Report to HSE submitted September 2007.
5. McNamee R, Carder M, Money A, Agius R. Time trends in the incidence of work-related disease in the UK, 1996-2007: estimation from ODIN/THOR surveillance data. Report to HSE submitted September 2008.
6. Carder M, McNamee R, Hussey L, Money A and Agius R. Time trends in the incidence of work-related disease in the UK, 1996-2008: estimation from THOR surveillance data. Report to HSE submitted September 2009.
7. Carder M, McNamee R, Hussey L, Money A and Agius R. Time trends in the incidence of work-related disease in the UK, 1996-2009: estimation from THOR surveillance data. Report to HSE submitted September 2010.
8. Carder M, McNamee R, Holland F, Hussey L, Money A and Agius R. Time trends in the incidence of work-related disease in the UK, 1996-2010: estimation from THOR surveillance data. Report to HSE submitted September 2011.
9. Carder M, McNamee R, Hussey L, Turner S and Agius R. Time trends in the incidence of work-related disease in the UK, 1996-2011: estimation from THOR surveillance data. Report to HSE submitted September 2012.
10. Carder M, McNamee R, Hussey L, and Agius R. Time trends in the incidence of work-related disease in the UK, 1996-2012: estimation from THOR surveillance data. Report to HSE submitted September 2013.
11. Carder M, McNamee R, Hussey L, and Agius R. Time trends in the incidence of work-related disease in the UK, 1996-2013: estimation from THOR surveillance data. Report to HSE submitted September 2014.
12. McNamee R, Carder M, Money A and Agius R. Time trends in the incidence of work-related disease in the UK, 1996-2006: estimation from ODIN/THOR

surveillance data. Supplement 1: Analysis of 'fatigue. Report to HSE submitted October 2007

13. Holland F and McNamee R. Work package 1: Modelling of zeros and non-response with membership time. Report submitted to HSE in January 2012
14. Holland F and McNamee R. Work package 2: Analysis of zero-inflated count data for EPIDERM, OPRA, THOR-GP and SWORD. Report submitted to HSE in July 2011
15. Holland F, McNamee R and Hodgson J. Summary of statistical work packages applied to THOR surveillance data. Report submitted to HSE in March 2012
16. Gittins M, McNamee R, Carter L and Holland F. Voluntary reporting schemes and the influence of excess zeroes on estimating true incidence. Submitted to The European Journal of Epidemiology.
17. Cherry, N, Meyer, J.D, Adisesh, A, Brooke, R, Owen-Smith, V, Swales, C & Beck, M.H. Surveillance of occupational skin disease: EPIDERM and OPRA. *Br J Derm*, 2000; 142: 1128-1134
18. Meredith SK, Taylor VM, McDonald JC. Occupational respiratory disease in the United Kingdom 1989: a report to the British Thoracic Society and the Society of Occupational Medicine by the SWORD project group. *Br J Ind Med* 1991; 48(5):292-298.
19. Hussey L. Calculating THOR-GP incidence rates from core and sample reported data. Report submitted to HSE, 2nd May 2013.
20. Office for National Statistics. Labour Force Survey 1996-2013: The Stationery Office, 2014
21. Firth D and De Menezes RX. Quasi-variances. *Biometrika* (2004); 91 (1): 65-80.
22. Turner S, McNamee R, Wilkinson SM, Agius R, Carder M, Stocks J. Evaluating interventions aimed at reducing occupational exposure to latex and rubber glove allergens. *Occupational and Environmental Medicine* 2012; 69(12):925-31.
23. Stocks SJ, McNamee R, Turner S et al. Has European Union legislation to reduce exposure to chromate in cement been effective in reducing the incidence of allergic contact dermatitis attributed to chromate in the UK? *Occup Environ Med* 2012;69:150-152.
24. Kwok C, Money A, Carder M, Turner S, Agius R, Orton D, and Wilkinson M. Occupational disease in Beauticians reported to The Health and Occupation Research (THOR) network from 1996 to 2011. *Clinical and Experimental Dermatology*, 2014; 39 (5): 590-595

25. Stocks SJ, McNamee R, Turner S, Carder M, Agius RM (2015) The impact of national level interventions to improve hygiene on the incidence of irritant contact dermatitis in healthcare workers: changes in incidence from 1996-2012 and interrupted times series analysis. *Br J Dermatol*. [Full text link doi: 10.1111/bjd.13719](https://doi.org/10.1111/bjd.13719). [Epub ahead of print]
26. Urwin R, Warburton K, Carder M, Turner S, Agius R, Wilkinson SM. (2015) Methylchloroisothiazolinone & methylisothiazolinone contact allergy: an occupational perspective. *Contact Dermatitis*, 72(6):381-6
27. Stock SJ *et al*. Trends in incidence of occupational asthma, contact dermatitis, noise-induced hearing loss, carpal tunnel syndrome and upper limb musculoskeletal disorders in European countries from 2000 to 2012 *Occup Environ Med*, 72:294-303
28. Health and Safety Executive. Self-reported work-related illness (SWI) and workplace injuries (LFS). Available at: <http://www.hse.gov.uk/statistics/publications/swi.htm> (last accessed August 2015).
29. Health and Safety Executive. Self-reported work-related illness (SWI) and workplace injuries: results from the Labour Force Survey (LFS). Incidence (3 year average) for people working in the last 12 months (Table SWIT6W12 3YR). Available at: <http://www.hse.gov.uk/statistics/lfs/index.htm> (last accessed August 2015).
30. Stocks SJ, McNamee R, Turner S, Carder M, Agius RM. Assessing the impact of national level interventions on workplace respiratory disease in the UK: part 1 - changes in workplace exposure legislation and market forces. *Occup Environ Med*, 2013; 70:476-482
31. Stocks SJ, McNamee R, Turner S, Carder M, Agius RM. Assessing the impact of national level interventions on workplace respiratory disease in the UK: part 2 - regulatory activity by the Health and Safety Executive. *Occup Environ Med*, 2013; 70:483-490.
32. Peto J, Hodgson J, Matthews F, Jones J (1995). Continuing increase in mesothelioma mortality in Britain. *The Lancet* 1995; 345:535-539.
33. Health and Safety Executive. Deaths from asbestos-related and other occupational lung diseases. Table MESO01: Death certificates mentioning mesothelioma, 1968-2013. Available at: <http://www.hse.gov.uk/statistics/tables/index.htm> (last accessed August 2015).
34. Pleural plaques litigation, House of Lords judgement. Available at: <http://www.publications.parliament.uk/pa/ld200607/ldjudgmt/jd071017/johns-1.htm> (last accessed August 2015).

APPENDIX A SUMMARY OF REPORTER FATIGUE INVESTIGATIONS

- 1) The probability of a zero return as a function of membership time:** the percentage increase, per year of membership, in the *odds* of a returned card having zero cases was estimated. These analyses were initially carried out for SWORD (1999-2004) and EPIDERM (1996-2004) and subsequently for THOR-GP (June 2005-2008). Separate analyses were carried out for 'core' and 'sample' reporters (except for THOR-GP, which was exclusively core reporting during this period). These analyses sought to separate the true trend with calendar time from a trend with membership time (used as a proxy for fatigue). Membership time was included as a covariate in the usual model which also included calendar time, season, and whether or not it was the first return. **Results:** Results were inconclusive due to wide confidence intervals caused by high collinearity between membership time and calendar time, especially for EPIDERM, SWORD and THOR-GP core reporters. There was some evidence for EPIDERM sample reporters that blank returns increased as a function of membership time (by 6% per year) but not for SWORD sample.
- 2) Calendar time trends in incidence adjusted for membership time:** The results of the analyses described in 1) suggested it might be possible to separate out the effects of calendar time and membership time for sample reporters. Therefore, the percentage change in incidence of total cases (EPIDERM 1996-2004, SWORD 1999-2004), 'adjusted' for an independent effect of membership time on incidence was estimated. Variables included in the MLM were 'calendar time', 'membership time', season, and first report. **Results:** Results suggested evidence of fatigue for EPIDERM sample reporters but not for SWORD sample reporters. On including 'membership time' in the models, the estimated annual change in incidence of cases reported to EPIDERM became -0.4% (95% CIs: -6.5, 6.2) instead of -3.2% whilst for SWORD it showed little change from -7.3% (95% CIs: -11.8, -2.7) to 7.1% (95% CIs: -12.0, -2.0).
- 3) Descriptive analysis using the FATCATS/CALCATS approach:** i.e. zero return rates broken down simultaneously by categories of membership time (2 year intervals) (FATCATS) and calendar time (2 year intervals) (CALCATS). This was initially undertaken for EPIDERM (1996-2006) and SWORD (1999-2006), and subsequently for THOR-GP (June 2005-2008). **Results:** EPIDERM and SWORD core: little evidence that for any given calendar period the proportion of zero returns increased with membership time or that for any given membership period the proportion of zero returns increased with calendar time. EPIDERM and SWORD sample: some evidence of the former but not of the latter phenomenon. THOR-GP core: little evidence of the former but some evidence of the latter phenomenon.

- 4) **GEE (generalised estimating equations) modelling on zero returns in relation to time:** The GEE modelling approach is an alternative to the random effects (RE) approach. It was used as a sensitivity analysis – to see if consistent with the results from RE approach. Zero return rates were modelled as a function of membership time, with adjustment for calendar time. Membership time was included in the model as either a continuous variable (years) or categorised (2 year intervals). Analyses were carried out on core and sample reporters combined (EPIDERM 1996-2006 and SWORD 1999-2006).

Results: Results suggested an increase in zero cases of 4% and 2% per membership year (EPIDERM and SWORD, respectively) but these trends were not statistically significant (EPIDERM $p=0.08$, SWORD $p=0.20$). In models where membership time was categorised, the odds ratios for all membership categories were higher than 1 (the reference year was <2 years membership) and seemed to settle around 1.3% after 6 years membership for EPIDERM whilst for SWORD there was no suggestion of an increase with membership time.

- 5) **Estimation of calendar time trends in incidence rates with membership restrictions:** The percentage change in incidence of WRI was estimated 'as usual' using the methodology described under Section 2.4 but reporters were categorised by membership time (2 year intervals) and separate analyses were carried out for each group. Analyses were carried out for core and sample reporters combined (EPIDERM 1996-2006, SWORD 1999-2006).

Results: The trends estimates suggested that there was some evidence that EPIDERM reporters, but not SWORD reporters, in the longer membership categories might be more influenced by fatigue (manifesting as an increase in zeros).

- 6) **Modelling of zeros and non-response with membership time:** Longitudinal logistic GEE and RE models were fitted to investigate the relationship between non-response and zero response with membership time i.e. whether the probability of either type of response changes as membership time increases, and whether one type of response is more likely than the other (and whether this changes with membership time).

Results: EPIDERM sample: there was strong evidence that both non-returns and zero returns (given a return) increased with membership time; the estimated odds were 13% and 7%, respectively. The conditional probability of a zero (i.e., given a zero case or non-return) declined over time (by 9% per membership year); we would expect this to decline if non-response increased more rapidly than zero returns. For the other reporters/schemes the estimated odds of non-response, zero response, and the conditional probability of a zero were EPIDERM core: 31%, 7% and 21%, respectively; SWORD sample: 17%, 4% and 14%. SWORD core: 33%, 7% and 18%, respectively.

All these analyses were conducted on total cases for each scheme. The implicit assumption is that fatigue was a general phenomenon affecting the reports as a whole for a given reporter and is not specific to a diagnostic group.

Table A1 Evidence of fatigue as exhibited by an increase in zero returns over time

Analyses	EPIDERM		SWORD		THOR-GP	
	Core	Sample	Core	Sample	Core	Sample
1*	/	Yes	/	No	/	/
2	/	Yes	/	No	/	/
3	No	Yes?	No	Yes?	Yes?	/
4**	Yes?		No		/	/
5	Yes?		No		/	/
6	Yes?	Yes	Yes?	Yes?	/	/

*It was not possible to separate out the effect of calendar time and membership time due to high collinearity between the two variables

**Analyses for SWORD and EPIDERM were on all reporters combined. This analysis was not repeated for THOR-GP

APPENDIX B DESCRIPTIVE ANALYSES

Table B1 Reporting activity of reporters in EPIDERM, 1996-2014

	CORE	SAMPLE
Total reporters ever in 1996-2014	57	392
Total active^a reporters in 1996-2014	55	362
Response rate^{**}	86%	75%
% of returns that are blank	17%	62%
Number of reporters who responded at least once but never returned a case	1	113
Number of reporters who have never responded	2	30

^a Active reporter is someone who returns a card

^b Response rate = cards returned/cards sent out

Figure B1 Number of reporters in EPIDERM by year and reporter type, 1996-2014






Figure B2 Response rates (cards returned/cards sent out) per year

a) All reporters



b) Core reporters



c) Sample reporters




Figure B3 Number of active reporters per month – EPIDERM, 1996-2014






Figure B4 Cases per active reporter per month – EPIDERM, 1996-2014


a) Total cases




b) Contact dermatitis



c) Contact urticaria (note scale change)



d) Neoplasia



e) Other skin (other than contact dermatitis) (note scale change)




Table B2 Cases reported per month by disease category and type of reporter, EPIDERM, 1996-2014

	Statistic	All Reporters				Core reporters				Sample reporters			
		Min	Max	SD		Min	Max	SD		Min	Max	SD	
Disease group All cases	Total active reporters ever in 1996-2014	392				55				362			
	Mean no. of active ^a reporters per month	30.22	16	42	6.27	19.76	9	26	4.06	10.46	3	20	3.18
	Total cases	18438				16239				2199			
	Mean cases per month	80.87	26	148	28.59	71.22	21	147	26.87	9.64	0	33	6.38
	Mean cases per active reporter per month	2.63	1.12	5.92	0.68	3.53	1.39	7.74	0.97	0.93	0	4.17	0.60
Contact dermatitis (CD)	Total cases	15113				13601				1512			
	Mean cases per month	66.29	22	122	23.14	59.65	20	121	21.87	6.63	0	23	4.60
	Mean cases per active reporter per month	2.17	0.97	4.88	0.58	2.98	1.33	6.37	0.81	0.64	0	2.83	0.44
Allergic CD	Total cases	5600				4949				651			
	Mean cases per month	24.56	4	58	11.10	21.71	3	54	10.16	2.86	0	12	2.64
	Mean cases per active reporter per month	0.79	0.21	1.66	0.29	1.07	0.23	2.44	0.40	0.27	0	1.83	0.26
Irritant CD	Total cases	6738				6179				559			
	Mean cases per month	29.55	8	58	10.65	27.10	6	58	10.41	2.45	0	12	2.16
	Mean cases per active reporter per month	0.98	0.41	2.32	0.31	1.36	0.50	3.05	0.43	0.24	0	1	0.21
Mixed CD	Total cases	2397				2195				202			
	Mean cases per month	10.51	1	27	5.07	9.63	1	25	4.93	0.89	0	5	1.11
	Mean cases per active reporter per month	0.35	0.05	0.92	0.15	0.48	0.05	1.21	0.22	0.09	0	0.67	0.12
Other ^b cases	Total cases	3751				3039				712			
	Mean cases per month	16.45	1	39	8.82	13.33	0	33	7.84	3.12	0	20	3.40

		All Reporters				Core reporters				Sample reporters			
Statistic			Min	Max	SD		Min	Max	SD		Min	Max	SD
Contact urticaria	Mean cases per active reporter per month	0.52	0.05	1.16	0.24	0.63	0	1.78	0.34	0.30	0	2.22	0.33
	Total cases	847				800				47			
	Mean cases per month	3.71	0	15	2.91	3.51	0	14	2.84	0.21	0	3	0.50
	Mean cases per active reporter per month	0.12	0	0.42	0.08	0.17	0	0.78	0.13	0.02	0	0.33	0.05
Neoplasia	Total cases	2221				1687				534			
	Mean cases per month	9.74	0	28	5.98	7.40	0	20	4.94	2.34	0	19	3.09
	Mean cases per active reporter per month	0.31	0	0.83	0.17	0.35	0	1.05	0.22	0.23	0	2.11	0.30

^aActive reporter is someone who returns a card

^bother than contact dermatitis

Table B3 Reporting activity of reporters in SWORD, 1999-2014

	CORE	SAMPLE
Total reporters ever in 1999-2014	48	813
Total active^a reporters in 1999-2014	44	765
Response rate^b	83%	73%
% of returns that are zero returns (i.e. no cases to report)	29%	72%
Number of reporters who responded at least once but never returned a case	1	251
Number of reporters who have never responded	4	48

^a Active reporter is someone who returns a card

^b Response rate = cards returned/cards sent out

Figure B5 Number of reporters in SWORD by year and reporter type






Figure B6 Response rates (cards returned/cards sent out) per year

a) All reporters



b) Core reporters



c) Sample reporters




Figure B7 Number of active reporters per month – SWORD






Figure B8 Cases per active reporter per month – SWORD


a) Total cases




b) Asthma (note scale change)




c) Mesothelioma



d) Non-malignant pleural disease (note scale change)



e) Pneumoconiosis (note scale change)



f) Other (than those specified above) respiratory disease




Table B4 Cases reported per month by disease category and type of reporter, SWORD, 1999-2014

		Statistic	All Reporters			Core reporters			Sample reporters					
			Min	Max	SD	Min	Max	SD	Min	Max	SD			
Disease group	All cases	Total active reporters ever in 1999-2014	789			44			765					
		Mean no. of active ^a reporters per month	43.30	22	59	8.65	15.93	7	24	4.42	27.36	13	38	5.02
		Total cases	12304			9754			2550					
		Mean cases per month	64.08	22	132	25.45	50.80	12	112	23.70	13.28	3	35	6.28
		Mean cases per active reporter per month	1.44	0.66	2.69	0.38	3.09	1.40	5.78	0.89	0.48	0.09	1.10	0.20
Asthma		Total cases	2362			2103			259					
		Mean cases per month	12.30	2	42	6.68	10.95	1	42	6.06	1.35	0	9	1.44
		Mean cases per active reporter per month	0.28	0.07	0.76	0.12	0.68	0.14	2.33	0.29	0.05	0	0.28	0.05
Mesothelioma		Total cases	2392			1560			832					
		Mean cases per month	12.46	0	34	6.78	8.13	0	27	5.78	4.33	0	11	2.65
		Mean cases per active reporter per month	0.28	0.00	0.67	0.12	0.47	0	1.69	0.28	0.16	0	0.45	0.09
Non-malignant plaques	pleural	Total cases	5265			4303			962					
		Mean cases per month	27.42	3	60	12.63	22.41	2	59	12.25	5.01	0	17	3.47
		Mean cases per active reporter per month	0.61	0.10	1.25	0.21	1.34	0.20	2.84	0.51	0.18	0	0.65	0.12
Pneumoconiosis		Total cases	1054			853			201					
		Mean cases per month	5.49	0	16	2.67	4.44	0	13	2.36	1.05	0	5	1.18

Statistic	All Reporters			Core reporters			Sample reporters					
	Min	Max	SD	Min	Max	SD	Min	Max	SD			
Mean cases per active reporter per month	0.13	0	0.35	0.06	0.29	0	1	0.17	0.04	0	0.20	0.04
Other cases ^b												
Total cases	1703				1360				343			
Mean cases per month	8.87	1	33	4.75	7.08	1	28	4.31	1.79	0	13	1.83
Mean cases per active reporter per month	0.20	0.02	0.60	0.10	0.46	0.05	1.56	0.26	0.06	0	0.45	0.06

^aActive reporter is someone who returns a card

^bOther than those specified above i.e SWORD categories: inhalation accidents, allergic alveolitis, bronchitis/emphysema, infectious disease, lung cancer and 'other' (the latter includes rhinitis). NOTE: A case may have more than one diagnosis

Table B5 Reporting activity of reporters in THOR-GP, 2006-2014

	CORE^a	SAMPLE
Total reporters ever in 2006-2014	442	313
Total active^b reporters in 2006-2014	332	262
Response rate^c	58%	70%
% of returns that are zero returns (i.e. no cases to report)	60%	39%
Number of reporters who responded at least once but never returned a case	46	53
Number of reporters who have never responded	110	51

^aCore reporting stopped in 2010

^bActive reporter is someone who returns a card

^cResponse rate = cards returned/cards sent out

Figure B9 Number of reporters in THOR-GP by year and reporter type






Figure B10 Response rates (cards returned/cards sent out) per year

a) All reporters



b) Core reporters



c) Sample reporters




Figure B11 Number of active reporters per month – THOR-GP






Figure B12 Cases per active reporter per month – THOR-GP


a) Total cases




b) Total skin (note scale change)




c) Contact dermatitis




d) Total musculoskeletal (note scale change)




e) Upper limb (note scale change)




f) Spine/back




g) Lower limb (note scale change)



h) Total mental ill-health (note scale change)



i) Anxiety and depression



j) Other work stress




Table B6 Cases reported per month by disease category and type of reporter, THOR-GP, 2006-2014

	Statistic	All Reporters				Core reporters				Sample reporters				SD
			Min	Max	SD		Min	Max	SD		Min	Max		
Disease group All cases	Total active reporters ever in 2006-2014	423				332				262				
	Mean no. of active ^a reporters per month	83.79	7	187	72.29	136.23	31	185	53.33	9.09	0	19	6.50	
	Total cases	6331				5115				1215				
	Mean cases per month	58.62	3	190	45.38	85.25	11	190	42.39	12.66	0	37	10.14	
	Mean cases per active reporter per month	0.94	0.35	3.70	0.48	0.62	0.33	1.44	0.20	1.54	0	7	1.25	
All skin	Total cases	613				508				105				
	Mean cases per month	5.68	0	25	5.21	8.47	0	25	5.06	1.09	0	6	1.49	
	Mean cases per active reporter per month	0.11	0	1.88	0.21	0.06	0	0.17	0.03	0.16	0	2	0.29	
Contact dermatitis	Total cases	472				400				72				
	Mean cases per month	4.37	0	21	4.27	6.67	0	21	4.25	0.75	0	6	1.13	
	Mean cases per active reporter per month	0.07	0	0.43	0.08	0.05	0	0.14	0.03	0.09	0	1	0.18	
All musculoskeletal	Total cases	3295				2713				582				
	Mean cases per month	30.51	1	106	25.46	45.22	5	106	24.53	6.06	0	17	5.08	
	Mean cases per active reporter per month	0.45	0	1.30	0.23	0.32	0.15	0.77	0.12	0.75	0	5	0.76	
Upper limb ^b	Total cases	1552				1304				248				
	Mean cases per month	14.37	1	52	12.52	21.73	1	52	11.87	2.58	0	8	2.22	
	Mean cases per active reporter per month	0.21	0.07	0.94	0.12	0.15	0.03	0.44	0.07	0.33	0	2	0.34	
Spine/back ^c	Total cases	1250				1007				243				

	Statistic	All Reporters			Core reporters			Sample reporters			SD	
		Min	Max	SD	Min	Max	SD	Min	Max	SD	Min	Max
	Mean cases per month	11.57	0	49	10.21	16.78	0	49	10.67	2.53	0	10
	Mean cases per active reporter per month	0.18	0.00	0.83	0.13	0.11	0	0.29	0.05	0.30	0	2
Lower limb ^d	Total cases	452				356				94		
	Mean cases per month	4.19	0	14	3.53	5.93	0	14	3.52	0.98	0	5
	Mean cases per active reporter per month	0.08	0.00	0.44	0.08	0.04	0	0.14	0.02	0.12	0	2
All mental ill-health	Total cases	2044				1604				453		
	Mean cases per month	18.93	1	56	13.99	26.73	3	56	13.03	4.72	0	18
	Mean cases per active reporter per month	0.33	0.00	1.50	0.22	0.20	0.08	0.44	0.07	0.51	0	3
Anxiety/depression	Total cases	890				700				180		
	Mean cases per month	8.24	0	26	6.63	11.67	1	26	6.33	1.88	0	11
	Mean cases per active reporter per month	0.14	0.00	1.10	0.15	0.09	0.02	0.21	0.04	0.18	0	2
Other work stress	Total cases	1347				1051				302		
	Mean cases per month	12.47	0	38	9.44	17.52	0	38	9.26	3.15	0	13
	Mean cases per active reporter per month	0.22	0.00	0.80	0.15	0.13	0	0.29	0.05	0.35	0	1

^aActive reporter is someone who returns a card

^bUpper limb = hand/wrist/arm, elbow and shoulder

^cSpine/back = neck/thoracic spine and lumbar spine/trunk

^dLower limb = ankle/knee/foot

NOTE: A case may have more than one diagnosis