

# National Confidential Inquiry

into Suicide and Safety  
in Mental Health

**Annual Report 2025:**  
**Additional online data tables**  
UK wide data 2012-2022\*

\*Findings include data from Jersey

## CONTENTS

A	PATIENT SUICIDE IN THE UK .....	3
B	THEMED TOPICS.....	4
	SUICIDE IN MENTAL HEALTH PATIENTS WITH BIPOLAR DISORDER .....	4
	SUICIDE FOLLOWING MISSED CONTACT .....	6
	SUICIDE FOLLOWING NON-ADHERENCE WITH MEDICATION .....	8
	SUICIDE AND RECENT (<3 MONTHS) BEREAVEMENT .....	10

## A PATIENT SUICIDE IN THE UK

**Table 1: Patient suicide: numbers by year and UK country**

	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>England</b>	1,379	1,321	1,241	1,270	1,258	1,256	1,415	1,350*	1,308*	1,327*	1,339*
<b>Northern Ireland<sup>†</sup></b>	78	77	61	43	54	52	67	83*	75*	71*	74*
<b>Scotland</b>	264	267	223	226	226	218	229	209	217*	210*	221*
<b>Wales</b>	96	101	63	79	64	85	85	85	90*	85*	83*
<b>Total</b>	<b>1,817</b>	<b>1,766</b>	<b>1,588</b>	<b>1,618</b>	<b>1,602</b>	<b>1,611</b>	<b>1,796</b>	<b>1,727*</b>	<b>1,690*</b>	<b>1,693*</b>	<b>1,717*</b>

<sup>†</sup>In Northern Ireland data prior to 2015 are not directly comparable with those in 2015 onwards due to a [review of suicide statistics by the Northern Ireland Statistics Agency \(NISRA\)](#)

\*Figures include estimates based on late notifications.

Data from Jersey are not shown due to the low number of patient deaths by suicide.

**Table 2: Patient suicide: number of in-patient suicides by year and UK country**

	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
<b>England</b>	93	97	77	90	66	62	71	66*	58*	57*	57*
<b>Northern Ireland</b>	4	3	<3	<3	<3	<3	<3	<3	<3	<3	<3
<b>Scotland</b>	19	16	12	20	14	6	13	8	9*	11*	11*
<b>Wales</b>	7	4	4	5	5	3	4	8	<3	6*	6*
<b>Total UK</b>	<b>123</b>	<b>120</b>	<b>93</b>	<b>115</b>	<b>85</b>	<b>71</b>	<b>88</b>	<b>82*</b>	<b>67*</b>	<b>74*</b>	<b>74*</b>

\*Figures include estimates based on late notifications.

Data from Jersey are not shown due to the low number of patient deaths by suicide.

## B THEMED TOPICS

### SUICIDE IN MENTAL HEALTH PATIENTS WITH BIPOLAR DISORDER (UK AND JERSEY, 2012-2022)

**Table 3: Socio-demographic characteristics of mental health patients with bipolar disorder compared to all other patients**

	Patients with bipolar disorder		All other patients
	Total=1,491 <sup>†</sup>		Total=15,794
	N	%	%
Age: median (range)	48 (17-90) △		46 (11-100)
Age group:			
Under 25	68	5% ▽	10%
25-44	476	34% ▽	37%
45-64	700	50% △	39%
≥65	170	12% ▽	14%
Female <sup>†</sup>	694	47% △	34%
Not currently married/co-habiting	972	73%	73%
Living alone	695	52% △	47%
Unemployed	653	49%	47%
LGBT+	38	6%	5%
Ethnic minority group	114	8%	7%
Homeless	21	2%	2%

△ ▽ = significantly (p<0.05) higher or lower than patients without a primary diagnosis of bipolar disorder

<sup>†</sup> includes estimated figures in 2020-2022

**Table 4: Behavioural characteristics and adverse life events of mental health patients with bipolar disorder compared to all other patients**

	Patients with bipolar disorder		Other patients
	Total=1,491 <sup>†</sup>		Total=15,794
	N	%	%
<b><i>Behavioural:</i></b>			
History of self-harm	823	61%	62%
History of violence	250	19%	19%
History of alcohol misuse	602	45%	47%
History of drug misuse	479	35%	38%
Suicide-related internet use	56	6% ▽	8%
<b><i>Recent (&lt;3 months) adverse events:</i></b>			
Serious financial problems	207	20% △	16%
Bereavement	85	7%	8%
Any life event	594	45% ▽	50%

△ ▽ = significantly (p<0.05) higher or lower than patients without a primary diagnosis of bipolar disorder

<sup>†</sup> includes estimated figures in 2020-2022

**Table 5: Clinical and service contact characteristics of mental health patients with bipolar disorder compared to all other patients**

<b>Characteristic</b>	<b>Patients with bipolar disorder</b>		<b>Other patients</b>
	<b>Total=1,491<sup>†</sup></b>		<b>Total=15,794</b>
	<b>N</b>	<b>%</b>	<b>%</b>
<b>Priority groups:</b>			
In-patients	77	5%	5%
Died within 3 months of discharge	210	16% △	13%
Under CRHT services	187	14%	13%
Missed last contact with services	280	21%	23%
Non-adherence with medication	201	15% △	12%
<b>Clinical:</b>			
Short (<12 months) history of illness	99	7% ▽	22%
Any comorbid mental illness	730	52% ▽	55%
Any physical illness	306	23% ▽	26%
<b>Service contact:</b>			
Recent (<1 week) contact	751	53% △	45%
Short-term risk: low or none	975	81%	81%
Long-term risk: low or none	636	56%	57%

△ ▽ = significantly (p<0.05) higher or lower than patients without a primary diagnosis of bipolar disorder

<sup>†</sup> includes estimated figures in 2020-2022

CRHT = crisis resolution/home treatment

## SUICIDE FOLLOWING MISSED CONTACT (UK AND JERSEY, 2012-2022)

**Table 6: Socio-demographic characteristics of mental health patients who had missed their last service contact compared to all other patients**

	Missed contact		All other patients
	Total=3,817 <sup>†</sup>		Total=12,487
	N	%	%
Age: median (range)	43 (11-94) ▽		47 (12-100)
Age group:			
Under 25	375	10% △	9%
25-44	1,586	44% △	35%
45-64	1,390	38% ▽	40%
≥65	270	7% ▽	16%
Male <sup>†</sup>	2,560	67% △	65%
Not currently married/co-habiting	2,646	80% △	70%
Living alone	1,883	56% △	45%
Unemployed	1,877	57% △	44%
LGBT+	95	6%	5%
Ethnic minority group	248	7%	7%
Homeless	101	3% △	2%

△ ▽ = significantly (p<0.05) higher or lower than other patients

<sup>†</sup> includes estimated figures in 2020-2022

**Table 7: Behavioural characteristics and adverse life events of mental health patients who had missed their last service contact compared to all other patients**

	Missed contact		Other patients
	Total=3,817 <sup>†</sup>		Total=12,487
	N	%	%
<b><i>Behavioural:</i></b>			
History of self-harm	2,236	65% △	60%
History of violence	771	24% △	17%
History of alcohol misuse	1,906	56% △	44%
History of drug misuse	1,687	49% △	34%
Suicide-related internet use	184	10% △	8%
<b><i>Recent (&lt;3 months) adverse events:</i></b>			
Serious financial problems	541	22% △	15%
Bereavement	290	10% △	8%
Any life event	1,609	51%	50%

△ ▽ = significantly (p<0.05) higher or lower than other patients

<sup>†</sup> includes estimated figures in 2020-2022

**Table 8: Clinical and service contact characteristics of mental health patients who had missed their last service contact compared to all other patients**

<i>Characteristic</i>	<b>Missed contact</b>		<b>Other patients</b>
	<b>Total=3,817<sup>†</sup></b>		<b>Total=12,487</b>
	<b>N</b>	<b>%</b>	<b>%</b>
<b><i>Clinical:</i></b>			
Primary diagnosis:			
Schizophrenia & other delusional disorders	542	15%	14%
Affective disorder (depression & bipolar disorder)	1,239	35% ▽	41%
Alcohol dependence/misuse	328	9% △	6%
Drug dependence/misuse	266	8% △	5%
Personality disorder	446	13% △	10%
Any comorbid mental illness	2,104	60% △	53%
Any physical illness	799	24% ▽	27%
Short (<12 months) history of illness	515	16% ▽	22%
<b><i>Priority groups:</i></b>			
Died within 3 months of discharge	449	13%	13%
Under CRHT services	488	14%	14%
Non-adherence with medication	571	18% △	11%
<b><i>Service contact:</i></b>			
Recent (<1 week) contact	1,219	34% ▽	47%
Short-term risk: low or none	2,412	81%	82%
Long-term risk: low or none	1,576	56% ▽	58%

△ ▽ = significantly (p<0.05) higher or lower than other patients

<sup>†</sup> includes estimated figures in 2020-2022

CRHT = crisis resolution/home treatment

## SUICIDE FOLLOWING NON-ADHERENCE WITH MEDICATION (UK AND JERSEY, 2012-2022)

**Table 9: Socio-demographic characteristics of mental health patients who were non-adherent with medication compared to all other patients**

	Non-adherent with medication		All other patients
	Total=1,998 <sup>†</sup>		Total=13,753
	N	%	%
Age: median (range)	44 (13-97) ▽		46 (11-100)
Age group:			
Under 25	197	10%	9%
25-44	766	40% △	36%
45-64	724	38%	40%
≥65	219	11% ▽	16%
Male <sup>†</sup>	1,298	65%	65%
Not currently married/co-habiting	1,386	76% △	72%
Living alone	877	48%	47%
Unemployed	937	52% △	46%
LGBT+	42	4%	5%
Ethnic minority group	158	9% △	7%
Homeless	49	3%	2%

△ ▽ = significantly (p<0.05) higher or lower than other patients

<sup>†</sup> includes estimated figures in 2020-2022

**Table 10: Behavioural characteristics and adverse life events of mental health patients who were non-adherent with medication compared to all other patients**

	Non-adherent with medication		Other patients
	Total=1,998 <sup>†</sup>		Total=13,753
	N	%	%
<b><i>Behavioural:</i></b>			
History of self-harm	1,140	62%	60%
History of violence	383	22% △	18%
History of alcohol misuse	855	48% △	45%
History of drug misuse	832	46% △	35%
Suicide-related internet use	134	11% △	7%
<b><i>Recent (&lt;3 months) adverse events:</i></b>			
Serious financial problems	278	19% △	16%
Bereavement	117	7% ▽	8%
Any life event	895	49%	50%

△ ▽ = significantly (p<0.05) higher or lower than other patients

<sup>†</sup> includes estimated figures in 2020-2022



**Table 11: Clinical and service contact characteristics of mental health patients who were non-adherent with medication compared to all other patients**

<i>Characteristic</i>	<b>Non-adherent with medication</b>		<b>Other patients</b>
	<b>Total=1,998<sup>†</sup></b>		<b>Total=13,753</b>
	<b>N</b>	<b>%</b>	<b>%</b>
<b><i>Clinical:</i></b>			
Primary diagnosis:			
Schizophrenia & other delusional disorders	458	24% △	14%
Affective disorder (depression & bipolar disorder)	722	38%	41%
Alcohol dependence/misuse	78	4% ▽	6%
Drug dependence/misuse	80	4%	5%
Personality disorder	198	11%	11%
Any comorbid mental illness	1,154	62% △	54%
Any physical illness	420	23% ▽	26%
Short (<12 months) history of illness	343	20% ▽	22%
<b><i>Priority groups:</i></b>			
In-patients	96	5%	6%
Died within 3 months of discharge	263	15%	14%
Under CRHT services	295	16%	14%
Missed last contact	571	32% △	20%
<b><i>Service contact:</i></b>			
Recent (<1 week) contact	987	52% △	49%
Short-term risk: low or none	1,191	72% ▽	82%
Long-term risk: low or none	707	45% ▽	57%

△ ▽ = significantly (p<0.05) higher or lower than other patients

<sup>†</sup> includes estimated figures in 2020-2022

CRHT = crisis resolution/home treatment

## SUICIDE AND RECENT (<3 MONTHS) BEREAVEMENT IN MENTAL HEALTH PATIENTS (UK AND JERSEY, 2012-2022)

**Table 9: Socio-demographic characteristics of recently (<3 months) bereaved mental health patients compared to all other patients**

	Recently bereaved		All other patients
	Total=1,312 <sup>†</sup>		Total=13,690
	N	%	%
Age: median (range)	49 (14-97) △		46 (11-100)
Age group:			
Under 25	104	8%	9%
25-44	391	32% ▽	37%
45-64	504	41%	40%
≥65	226	18% △	14%
Female <sup>†</sup>	526	40% △	35%
Not currently married/co-habiting	889	78% △	72%
Living alone	629	54% △	47%
Unemployed	519	46%	46%
LGBT+	39	5%	5%
Ethnic minority group	58	5% ▽	7%
Homeless	26	2%	2%

△ ▽ = significantly (p<0.05) higher or lower than other patients

<sup>†</sup> includes estimated figures in 2020-2022

**Table 10: Behavioural characteristics and adverse life events of recently (<3 months) bereaved mental health patients compared to all other patients**

	Recently bereaved		Other patients
	Total=1,312 <sup>†</sup>		Total=13,690
	N	%	%
<b><i>Behavioural:</i></b>			
History of self-harm	747	63%	60%
History of violence	182	16%	18%
History of alcohol misuse	539	47%	44%
History of drug misuse	402	34%	35%
Suicide-related internet use	60	8%	8%
<b><i>Recent (&lt;3 months) adverse events:</i></b>			
Serious financial problems	178	19% △	16%

△ ▽ = significantly (p<0.05) higher or lower than other patients

<sup>†</sup> includes estimated figures in 2020-2022

**Table 11: Clinical and service contact characteristics of recently (<3 months) bereaved mental health patients compared to all other patients**

<b>Characteristic</b>	<b>Recently bereaved</b>		<b>Other patients</b>
	<b>Total=1,312<sup>†</sup></b>		<b>Total=13,690</b>
	<b>N</b>	<b>%</b>	<b>%</b>
<b><i>Clinical:</i></b>			
Primary diagnosis:			
Schizophrenia & other delusional disorders	123	10% ▽	16%
Affective disorder (depression & bipolar disorder)	490	41%	41%
Alcohol dependence/misuse	81	7%	6%
Drug dependence/misuse	40	3%	5%
Personality disorder	143	12%	11%
Adjustment disorder	108	9%	5%
Any comorbid mental illness	720	61% △	54%
Any physical illness	320	29% △	25%
Short (<12 months) history of illness	276	26% △	22%
<b><i>Priority groups:</i></b>			
In-patients	64	5%	6%
Died within 3 months of discharge	155	13%	15%
Under CRHT services	199	17% △	15%
Missed last contact	290	26% △	21%
Non-adherent with medication	117	10% ▽	13%
<b><i>Service contact:</i></b>			
Recent (<1 week) contact	611	50%	51%
Short-term risk: low or none	788	75% ▽	81%
Long-term risk: low or none	506	52% ▽	56%

△ ▽ = significantly (p<0.05) higher or lower than other patients

<sup>†</sup> includes estimated figures in 2020-2022

CRHT = crisis resolution/home treatment