

Provisional patient suicide data (2012-2025)

Provisional numbers of patient suicide deaths in the UK and Jersey. Includes deaths occurring in 2012-2023 and provisional data for deaths occurring in 2024-2025.

About this data

NCISH data covers suicides by people in contact with mental health services in the 12 months before death (referred to as ‘patients’). Complete details of the NCISH methodology are provided [here](#).

Our latest [NCISH annual report](#) includes findings on patients who died by suicide in the baseline year of 2013 and the subsequent 10 years. 2023 is the most recent year for which comprehensive data are available.

Here we present the most up-to-date suicide data, including all data we currently hold for deaths that occurred between 2012 and 2025 in the UK and Jersey. The years 2024 and 2025 are incomplete and will be updated on a quarterly basis as more data become available.

The table below provides a summary of the key characteristics of patients who died by suicide. These data will be updated in January 2027. NCISH data for 2026 will be published from July 2027.

Characteristics of mental health patients who died by suicide (UK and Jersey, 2012-2025)

Characteristic	N = 21,566
	N (valid %)
Sociodemographic:	
Age: Median (min-max)	46 (11-100)
Male	14,035 (65%)
Female	7,531 (35%)
Living alone	9,476 (47%)
Unemployed	9,023 (47%)
Clinical:	
In-patient at time of suicide	1,131 (5%)
Under crisis resolution home treatment services	2,812 (14%)
Died within 3 months of discharge from in-patient care	2,680 (13%)
Primary diagnosis:	
Schizophrenia and other primary psychotic disorders	3,117 (15%)
Depressive illness	6,464 (31%)
Bipolar disorder	1,694 (8%)
Alcohol dependence/misuse	1,355 (6%)
Drug dependence/misuse	1,046 (5%)
Personality disorder	2,325 (11%)
Anxiety disorders (including PTSD)	1,585 (8%)
Other diagnosis*	2,654 (13%)
Any comorbid (additional) mental health diagnosis	11,526 (55%)
Behavioural:	
History of self-harm	12,477 (61%)
History of alcohol misuse	9,350 (46%)
History of drug misuse	7,712 (38%)
Risk:	
Short-term risk viewed as low or not present	14,276 (81%)

*other diagnoses include eating disorders, learning disability, conduct disorder, adjustment disorder, autism spectrum disorder, ADHD, somatisation disorder, organic disorder, drug induced psychosis, dementia and other specified.